

WEYL GROUPOIDS WITH AT MOST THREE OBJECTS

M. CUNTZ AND I. HECKENBERGER

ABSTRACT. We adapt the generalization of root systems of the second author and H. Yamane to the terminology of category theory. We introduce Cartan schemes, associated root systems and Weyl groupoids. After some preliminary general results, we completely classify all finite Weyl groupoids with at most three objects. The classification yields that there exist infinitely many “standard”, but only 9 “exceptional” cases.

Key words: Nichols algebra, reflection, root system, Weyl groupoid

1. INTRODUCTION

In the last decades, root systems and their generalizations have continuously led to many remarkable new results. In most cases the motivation was to understand the structure of some generalization of Lie algebras, for example Kac-Moody algebras or Lie superalgebras.

Following the plan of Andruskiewitsch and Schneider for a classification of pointed Hopf algebras [AS98],[AS00], a new type of root systems emerged [Hec06b]. These are fundamental invariants of Nichols algebras of diagonal type, and are crucial for the full classification of finite-dimensional Nichols algebras of diagonal type [Hec06a]. In [HY08] an axiomatic definition of a generalization of root systems was introduced, based on the main properties of the root systems of Nichols algebras of diagonal type. The class of these root systems includes properly the reduced root systems in the sense of Bourbaki [Bou68] and the root systems of Kac-Moody algebras [Kac90], but contains many exceptional cases. The reduced root systems of simple Lie superalgebras also fit naturally into the new framework [HY08]. The results in [AHS08] indicate that a large class of Nichols algebras of non-diagonal type, such as those of finite group type, presumably admits a root system satisfying the axioms given in [HY08].

I.H. is supported by the German Research Foundation (DFG) via a Heisenberg fellowship.

The main aspect of the novel root systems is, that one starts with a family of Cartan matrices instead of a single Cartan matrix. Consequently, the symmetry object is not a group but a groupoid, the so called *Weyl groupoid*.

There have also been many efforts to find “root systems” for complex reflection groups [BMM99], the main achievement being the cyclotomic Hecke algebras. In a connected groupoid, if one fixes an object a , then the morphisms from a to a form a group $\text{Hom}(a)$ which is isomorphic for all choices of a . It is easy to see that any finite group appears as $\text{Hom}(a)$ for some finite Weyl groupoid. So in particular, it will be interesting to investigate the root systems for which $\text{Hom}(a)$ is a complex reflection group.

Matsumoto’s theorem holds for Coxeter groupoids [HY08] and hence there will be many nice properties of Coxeter groups which may be translated to this new situation. However, there are some important differences, for example the exchange condition only holds in a weak version.

In this article, we focus on various different aspects of Weyl groupoids. To stress the naturality of the construction, we introduce new concepts for the definitions using the language of category theory. A Weyl groupoid is based on a set of Cartan matrices $\mathcal{C}$ called a *Cartan scheme*. For such a scheme $\mathcal{C}$, we define root systems of type $\mathcal{C}$ and their Weyl groupoid. We stay in the general setting and deduce many useful results about root systems, Cartan schemes and Weyl groupoids, extending the analysis in [HY08]. We discuss standard Cartan schemes and their Weyl groupoids: Regardless of the number of objects, these are defined with the help of a single Cartan matrix, and are closely related to the crystallographic Weyl groups. Then decompositions of Cartan schemes, root systems, and their Weyl groupoids are investigated and characterized. In Prop. 4.6 we prove that a finite root system for a given Cartan scheme is reducible if and only if the family of Cartan matrices is simultaneously decomposable, or equivalently, if one of the Cartan matrices of the family is decomposable.

Then we turn our attention to the case of finite root systems. The main theorems merge to the result:

Theorem 1.1. *Let $\mathcal{W}$ be the Weyl groupoid of a finite irreducible connected root system with at most 3 objects. Then one of the following holds:*

- (1) $\mathcal{W}$ is standard, i.e. all Cartan matrices are equal.
- (2) $\mathcal{W}$ is one of 9 exceptional Weyl groupoids.

The main tool in our proofs is Thm. 2.6, the proof of which is given in [HY08]. It states that $\mathcal{W}$ is generated by reflections and Coxeter relations. For details on which Cartan matrices actually yield standard Weyl groupoids and a description of the exceptional cases, see the theorems 5.4, 6.1, 6.3 and 6.5. As a consequence of our classification, we conclude in Rems. 5.5 and 6.2 that there exist root systems associated to some non-standard Cartan schemes which cannot be obtained as a root system of a finite-dimensional Nichols algebra of diagonal type.

There are many open questions left. It is conceivable that there are only finitely many non-standard irreducible connected Weyl groupoids for a fixed number of objects. Notice that there are infinitely many standard irreducible connected Weyl groupoids with two objects, but that all irreducible connected Weyl groupoids with three objects have rank less or equal four.

We use the symbols $\mathbb{N}$ and $\mathbb{N}_0$ for the set of positive and nonnegative integers, respectively.

We want to thank G. Malle for providing us with Ex. 3.2, and H.-J. Schneider for many interesting discussions on the subject and for his help in looking for a good terminology.

2. CARTAN SCHEMES, ROOT SYSTEMS, AND THEIR WEYL GROUPOIDS

First the generalization of root systems given in [HY08, Def. 2] is reformulated in terms of category theory.

Let I be a non-empty finite set and $\{\alpha_i \mid i \in I\}$ the standard basis of $\mathbb{Z}^I$. Recall from [Kac90, §1.1] that a generalized Cartan matrix $C = (c_{ij})_{i,j \in I}$ is a matrix in $\mathbb{Z}^{I \times I}$ such that

- (M1) $c_{ii} = 2$ and $c_{jk} \leq 0$ for all $i, j, k \in I$ with $j \neq k$,
- (M2) if $i, j \in I$ and $c_{ij} = 0$, then $c_{ji} = 0$.

Definition 2.1. Let A be a non-empty set, $\rho_i : A \rightarrow A$ a map for all $i \in I$, and $C^a = (c_{jk}^a)_{j,k \in I}$ a generalized Cartan matrix in $\mathbb{Z}^{I \times I}$ for all $a \in A$. The quadruple

$$\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$$

is called a *Cartan scheme* if

- (C1) $\rho_i^2 = \text{id}$ for all $i \in I$,
- (C2) $c_{ij}^a = c_{ij}^{\rho_i(a)}$ for all $a \in A$ and $i, j \in I$.

We say that $\mathcal{C}$ is *connected*, if the group $\langle \rho_i \mid i \in I \rangle \subset \text{Aut}(A)$ acts transitively on A , that is, if for all $a, b \in A$ with $a \neq b$ there exist

$n \in \mathbb{N}$, $a_1, a_2, \dots, a_n \in A$, and $i_1, i_2, \dots, i_{n-1} \in I$ such that

$$a_1 = a, \quad a_n = b, \quad a_{j+1} = \rho_{i_j}(a_j) \quad \text{for all } j = 1, \dots, n-1.$$

Two Cartan schemes $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ and $\mathcal{C}' = \mathcal{C}'(I', A', (\rho'_i)_{i \in I'}, (C'^a)_{a \in A'})$ are termed *equivalent*, if there are bijections $\varphi_0 : I \rightarrow I'$ and $\varphi_1 : A \rightarrow A'$ such that

$$\varphi_1(\rho_i(a)) = \rho'_{\varphi_0(i)}(\varphi_1(a)), \quad c_{\varphi_0(i)\varphi_0(j)}^{\varphi_1(a)} = c_{ij}^a$$

for all $i, j \in I$ and $a \in A$.

Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme. For all $i \in I$ and $a \in A$ define $\sigma_i^a \in \text{Aut}(\mathbb{Z}^I)$ by

$$(2.1) \quad \sigma_i^a(\alpha_j) = \alpha_j - c_{ij}^a \alpha_i \quad \text{for all } j \in I.$$

The *Weyl groupoid of $\mathcal{C}$* is the category $\mathcal{W}(\mathcal{C})$ such that $\text{Ob}(\mathcal{W}(\mathcal{C})) = A$ and the morphisms are generated by the maps $\sigma_i^a \in \text{Hom}(a, \rho_i(a))$ with $i \in I$, $a \in A$. Formally, for $a, b \in A$ the set $\text{Hom}(a, b)$ consists of the triples (b, f, a) , where

$$f = \sigma_{i_n}^{\rho_{i_{n-1}} \cdots \rho_{i_1}(a)} \cdots \sigma_{i_2}^{\rho_{i_1}(a)} \sigma_{i_1}^a$$

and $b = \rho_{i_n} \cdots \rho_{i_2} \rho_{i_1}(a)$ for some $n \in \mathbb{N}_0$ and $i_1, \dots, i_n \in I$. The composition is induced by the group structure of $\text{Aut}(\mathbb{Z}^I)$:

$$(a_3, f_2, a_2) \circ (a_2, f_1, a_1) = (a_3, f_2 f_1, a_1)$$

for all $(a_3, f_2, a_2), (a_2, f_1, a_1) \in \text{Hom}(\mathcal{W}(\mathcal{C}))$. By abuse of notation we will write $f \in \text{Hom}(a, b)$ instead of $(b, f, a) \in \text{Hom}(a, b)$.

The cardinality of I is termed the *rank of $\mathcal{W}(\mathcal{C})$* .

The Weyl groupoid $\mathcal{W}(\mathcal{C})$ of a Cartan scheme $\mathcal{C}$ is a groupoid. Indeed, (M1) implies that $\sigma_i^a \in \text{Aut}(\mathbb{Z}^I)$ is a reflection for all $i \in I$ and $a \in A$, and hence the inverse of $\sigma_i^a \in \text{Hom}(a, \rho_i(a))$ is $\sigma_i^{\rho_i(a)} \in \text{Hom}(\rho_i(a), a)$ by (C1) and (C2). Therefore each morphism of $\mathcal{W}(\mathcal{C})$ is an isomorphism.

If $\mathcal{C}$ and $\mathcal{C}'$ are equivalent Cartan schemes, then $\mathcal{W}(\mathcal{C})$ and $\mathcal{W}(\mathcal{C}')$ are isomorphic groupoids.

Recall that a groupoid G is called *connected*, if for each $a, b \in \text{Ob}(G)$ the class $\text{Hom}(a, b)$ is non-empty. Hence $\mathcal{W}(\mathcal{C})$ is a connected groupoid if and only if $\mathcal{C}$ is a connected Cartan scheme.

Definition 2.2. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme. For all $a \in A$ let $R^a \subset \mathbb{Z}^I$, and define $m_{i,j}^a = |R^a \cap (\mathbb{N}_0 \alpha_i + \mathbb{N}_0 \alpha_j)|$ for all $i, j \in I$ and $a \in A$. We say that

$$\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$$

is a *root system of type $\mathcal{C}$* , if it satisfies the following axioms.

- (R1) $R^a = R_+^a \cup -R_+^a$, where $R_+^a = R^a \cap \mathbb{N}_0^I$, for all $a \in A$.
- (R2) $R^a \cap \mathbb{Z}\alpha_i = \{\alpha_i, -\alpha_i\}$ for all $i \in I$, $a \in A$.
- (R3) $\sigma_i^a(R^a) = R^{\rho_i(a)}$ for all $i \in I$, $a \in A$.
- (R4) If $i, j \in I$ and $a \in A$ such that $i \neq j$ and $m_{i,j}^a$ is finite, then $(\rho_i \rho_j)^{m_{i,j}^a}(a) = a$.

If $\mathcal{R}$ is a root system of type $\mathcal{C}$, then we say that $\mathcal{W}(\mathcal{R}) = \mathcal{W}(\mathcal{C})$ is the *Weyl groupoid of $\mathcal{R}$* . Further, $\mathcal{R}$ is called *connected*, if $\mathcal{C}$ is a connected Cartan scheme. If $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ is a root system of type $\mathcal{C}$ and $\mathcal{R}' = \mathcal{R}'(\mathcal{C}', (R'^a)_{a \in A'})$ is a root system of type $\mathcal{C}'$, then we say that $\mathcal{R}$ and $\mathcal{R}'$ are *equivalent*, if $\mathcal{C}$ and $\mathcal{C}'$ are equivalent Cartan schemes given by maps $\varphi_0 : I \rightarrow I'$, $\varphi_1 : A \rightarrow A'$ as in Def. 2.1, and if the map $\varphi_0^* : \mathbb{Z}^I \rightarrow \mathbb{Z}^{I'}$ given by $\varphi_0^*(\alpha_i) = \alpha_{\varphi_0(i)}$ satisfies $\varphi_0^*(R^a) = R'^{\varphi_1(a)}$ for all $a \in A$.

Remark 2.3. (1) Reduced root systems with a fixed basis, see [Bou68, Ch. VI, §1.5], are examples of root systems of type $\mathcal{C}$ in the following way. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme, such that $A = \{a\}$ has only one element, and C^a is a Cartan matrix of finite type. Then $\rho_i = \text{id}$ for all $i \in I$. Let R^a be the reduced root system associated to C^a , where the basis $\{\alpha_i \mid i \in I\}$ of $\mathbb{Z}^I$ is identified with a basis of R^a . Then $\mathcal{R} = \mathcal{R}(\mathcal{C}, R^a)$ is a root system of type $\mathcal{C}$.

(2) In root systems 0 is never a root. The same holds for the sets R^a in root systems of type $\mathcal{C}$. Indeed, if $0 \in R^a$, then $0 \in R^a \cap \mathbb{Z}\alpha_i$ for all $i \in I$, and since I is non-empty, this is a contradiction to (R2).

(3) Let $\mathcal{C}$ be a Cartan scheme and $\mathcal{R}$ a root system of type $\mathcal{C}$. For $i, j \in I$ with $i \neq j$, (C1) and (R4) imply that the relations

$$(2.2) \quad (\rho_j \rho_i)^{m_{i,j}^a}(a) = a$$

hold for all $a \in A$. Further,

$$(2.3) \quad (\sigma_i \sigma_j)^{m_{i,j}^a} 1_a = (\sigma_j \sigma_i)^{m_{i,j}^a} 1_a = 1_a$$

for all $a \in A$ and $i, j \in I$ with $i \neq j$, see Thm. 2.6 below. Here 1_a is the identity of the object a , and we use the convention that upper indices referring to objects are neglected if they are uniquely determined by the context.

Remark 2.4. In [HY08, Def. 2] it is assumed that a root system $\mathcal{R}$ of type $\mathcal{C}$ is *connected*. We omit this axiom to have a definition which is compatible with passing to restrictions, see Def. 4.1. Note that a restriction of $\mathcal{R}$ is generally not connected, even if $\mathcal{R}$ is connected.

The following lemma states that in root systems of type $\mathcal{C}$ some axioms are redundant.

Lemma 2.5. *Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a quadruple, where A is a non-empty set, $\rho_i : A \rightarrow A$ is a map for all $i \in I$, and $C^a = (c_{ij}^a)_{i,j \in I} \in \mathbb{Z}^{I \times I}$ for all $a \in A$, such that $c_{ii}^a = 2$ for all $i \in I$, $a \in A$, and that (C1) holds. For all $a \in A$ let $R^a \subset \mathbb{Z}^I$ satisfying (R1)–(R4). Then $\mathcal{C}$ is a Cartan scheme and $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ is a root system of type $\mathcal{C}$.*

Proof. We have to prove that (M1), (C2), and (M2) hold for C^a for all $a \in A$. Let $a \in A$ and $j, k \in I$ with $j \neq k$. Then $\alpha_k \in R^a$ by (R2), hence $\sigma_j^a(\alpha_k) \in R_+^{\rho_j(a)}$ by (R1) and (R3). Therefore $c_{jk}^a \leq 0$ by Eq. (2.1). This proves (M1) for C^a .

Let now $a \in A$ and $i, j \in I$. Then

$$\begin{aligned} \sigma_i^{\rho_i(a)} \sigma_i^a(\alpha_j) &= \sigma_i^{\rho_i(a)}(\alpha_j - c_{ij}^a \alpha_i) \\ &= \alpha_j + (c_{ij}^a - c_{ij}^{\rho_i(a)}) \alpha_i \in R_+^{\rho_i^2(a)} = R_+^a \end{aligned}$$

by Eq. (2.1), (R1)–(R3), and (C1), and hence $c_{ij}^a \geq c_{ij}^{\rho_i(a)}$. Replacing a by $\rho_i(a)$, we obtain in the same way that $c_{ij}^a \leq c_{ij}^{\rho_i(a)}$. Hence (C2) holds. Assume now that $c_{ij}^a = 0$. Then $i \neq j$ by (M1). Further, Eq. (2.1), (C2), and relation $c_{ij}^a = 0$ imply that

$$\sigma_i^a \sigma_j^{\rho_j(a)}(\alpha_i) = \sigma_i^a(\alpha_i - c_{ji}^{\rho_j(a)} \alpha_j) = -\alpha_i - c_{ji}^a \alpha_j,$$

and $\sigma_i^a \sigma_j^{\rho_j(a)}(\alpha_i) \in -R_+^{\rho_i(a)}$ by (R1)–(R3). Since $c_{ji}^a \leq 0$ by (M1), this gives that $c_{ji}^a = 0$, hence (M2) is proven. $\square$

Recall the convention in Rem. 2.3(3). The Weyl groupoid of a root system of type $\mathcal{C}$ is a generalization of the notion of a Weyl group, as the following theorem shows.

Theorem 2.6. [HY08, Thm. 1] *Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme and $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ a root system of type $\mathcal{C}$. Let $\mathcal{W}$ be the abstract groupoid with $\text{Ob}(\mathcal{W}) = A$ such that $\text{Hom}(\mathcal{W})$ is generated by abstract morphisms $s_i^a \in \text{Hom}(a, \rho_i(a))$, where $i \in I$ and $a \in A$, satisfying the relations*

$$s_i s_i 1_a = 1_a, \quad (s_j s_k)^{m_{j,k}^a} 1_a = 1_a, \quad a \in A, i, j, k \in I, j \neq k.$$

Here 1_a is the identity of the object a , and $(s_j s_k)^\infty 1_a$ is understood to be 1_a . The functor $\mathcal{W} \rightarrow \mathcal{W}(\mathcal{R})$, which is the identity on the objects, and on the set of morphisms is given by $s_i^a \mapsto \sigma_i^a$ for all $i \in I$, $a \in A$, is an isomorphism of groupoids.

One says that $\mathcal{W}(\mathcal{R})$ is a *Coxeter groupoid*. Thus it makes sense to speak about the *length*

$$(2.4) \quad \ell(\omega) = \min\{m \in \mathbb{N}_0 \mid \omega = \sigma_{i_1} \cdots \sigma_{i_m} 1_a, i_1, \dots, i_m \in I\}$$

of a morphism $\omega \in \text{Hom}(a, b) \subset \text{Hom}(\mathcal{W}(\mathcal{R}))$. The most essential difference between Coxeter groupoids and Coxeter groups is the presence of several objects in the former.

Definition 2.7. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme. Let Γ be a nondirected graph, such that each edge is labelled by an element of I , and any two edges between two fixed vertices are labelled differently. Assume that there is a bijection φ from A to the set of vertices of Γ , and two vertices $\varphi(a), \varphi(b)$, where $a, b \in A$, are connected by an edge labelled by $i \in I$ if and only if $a \neq b$ and $\rho_i(a) = b$. The graph Γ is called the *object change diagram* of $\mathcal{C}$. If $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ is a root system of type $\mathcal{C}$, then we also say that Γ is the object change diagram of $\mathcal{R}$.

The object change diagram of a reduced root system is a single vertex without any edges. Other examples will appear in later sections. Note that the object change diagram of a Cartan scheme $\mathcal{C}$ is connected as a graph if and only if the Cartan scheme $\mathcal{C}$ is connected, or equivalently, if the Weyl groupoid $\mathcal{W}(\mathcal{C})$ is a connected groupoid.

Definition 2.8. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme and $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ a root system of type $\mathcal{C}$. For all $a \in A$ let $(R^a)^{\text{re}} = \{\omega(\alpha_i) \mid \omega \in \text{Hom}(b, a), b \in A, i \in I\}$, and call $(R^a)^{\text{re}}$ the *set of real roots* of a .

Note that by (R2) and (R3) we have $(R^a)^{\text{re}} \subset R^a$ for all $a \in A$. The sets of real roots are interesting for various reasons, one of them is the following.

Proposition 2.9. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme, and let $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ be a root system of type $\mathcal{C}$. Then $\mathcal{R}^{\text{re}} = \mathcal{R}^{\text{re}}(\mathcal{C}, ((R^a)^{\text{re}})_{a \in A})$ is a root system of type $\mathcal{C}$, and $\mathcal{W}(\mathcal{R}^{\text{re}}) = \mathcal{W}(\mathcal{R})$.

Proof. Since $\mathcal{C}$ is a Cartan scheme, it suffices to show that the sets $(R^a)^{\text{re}}$ satisfy axioms (R1)–(R4) for all $a \in A$. Let $a \in A$. Since $\omega \sigma_i^{\rho_i(b)}(\alpha_i) = -\omega(\alpha_i)$ for all $i \in I, b \in A$, and $\omega \in \text{Hom}(b, a)$, we obtain that $(R^a)^{\text{re}} = -(R^a)^{\text{re}}$. Let $(R^a)_+^{\text{re}} = (R^a)^{\text{re}} \cap \mathbb{N}_0^I$ and $(R^a)_-^{\text{re}} = (R^a)^{\text{re}} \cap -\mathbb{N}_0^I$. Then $(R^a)_-^{\text{re}} = -(-(R^a)^{\text{re}} \cap \mathbb{N}_0^I) = -(R^a)_+^{\text{re}}$, and hence (R1) implies that $(R^a)^{\text{re}} = (R^a)_+^{\text{re}} \cup -(R^a)_+^{\text{re}}$. Since $\alpha_i \in (R^a)^{\text{re}}$, $(R^a)^{\text{re}} \subset R^a$, and $(R^a)^{\text{re}} = -(R^a)^{\text{re}}$ by (R1), Axiom (R2) implies that $(R^a)^{\text{re}} \cap \mathbb{Z}\alpha_i =$

$\{\alpha_i, -\alpha_i\}$. Axiom (R3) holds for $\mathcal{R}^{\text{re}}$ by definition. Finally, if $m_{i,j}^a$ is finite for an $a \in A$ and elements $i, j \in I$ with $i \neq j$, then

$$R^a \cap (\mathbb{N}_0 \alpha_i + \mathbb{N}_0 \alpha_j) = (R^a)^{\text{re}} \cap (\mathbb{N}_0 \alpha_i + \mathbb{N}_0 \alpha_j)$$

by [HY08, Lemma 4]. Thus (R4) holds for $\mathcal{R}^{\text{re}}$, since it holds for $\mathcal{R}$. The equation $\mathcal{W}(\mathcal{R}^{\text{re}}) = \mathcal{W}(\mathcal{R})$ follows since $\mathcal{W}(\mathcal{R}^{\text{re}}) = \mathcal{W}(\mathcal{C}) = \mathcal{W}(\mathcal{R})$ by definition. $\square$

Now we discuss the finiteness of root systems of type $\mathcal{C}$.

Definition 2.10. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme and $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ a root system of type $\mathcal{C}$. We say that $\mathcal{R}$ is *finite* if R^a is finite for all $a \in A$.

The finiteness of $\mathcal{R}$ does not mean that $\mathcal{W}(\mathcal{R})$ is finite, since A may be infinite. But the following holds.

Lemma 2.11. *Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a connected Cartan scheme and $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ a root system of type $\mathcal{C}$. Then the following are equivalent.*

- (1) $\mathcal{R}$ is finite.
- (2) R^a is finite for at least one $a \in A$.
- (3) $\mathcal{R}^{\text{re}}$ is finite.
- (4) $\mathcal{W}(\mathcal{R})$ is finite.

Proof. (1) $\Rightarrow$ (2) is trivial.

(2) $\Rightarrow$ (1). Assume that $a \in A$ such that R^a is finite. Since $\mathcal{R}$ is connected, for each $b \in A$ there exists $\omega \in \text{Hom}(a, b)$. Then $R^b = \omega(R^a)$ by (R3), and hence R^b is finite for all $b \in A$.

(1) $\Rightarrow$ (4). Let $a, b \in A$. Since R^b is finite, R^a contains the standard basis of $\mathbb{Z}^I$ by (R2), and since $\omega(R^a) \subset R^b$ for all $\omega \in \text{Hom}(a, b)$, the set $\text{Hom}(a, b)$ is finite. Assume now that A is infinite, and let $a \in A$. The finiteness of R^a implies that there exist $b, c \in A$ and $f \in \text{Hom}(a, b)$, $g \in \text{Hom}(a, c)$ with $f \neq g$ and

$$Y := \{\alpha \in R_+^a \mid f(\alpha) \in R_+^b\} = \{\beta \in R_+^a \mid g(\beta) \in R_+^c\}.$$

Then $fg^{-1} \in \text{Hom}(c, b)$ and

$$\begin{aligned} fg^{-1}(g(Y)) &= f(Y) \subset R_+^b, \\ fg^{-1}(R_+^c \setminus g(Y)) &= f(-(R_+^a \setminus Y)) \subset R_+^b \end{aligned}$$

by (R1) and (R3). Therefore $fg^{-1}(R_+^c) \subset R_+^b$, and hence $b = c$ and $fg^{-1} = 1_b$ by [HY08, Lemma 8(iii)]. This is a contradiction to $f \neq g$, and hence A is finite. This proves (1) $\Rightarrow$ (4).

(4) $\Rightarrow$ (1). We prove that if $\mathcal{R}$ is infinite, then $\mathcal{W}(\mathcal{R})$ is infinite. For this we show by induction on m that for all $m \in \mathbb{N}$ there exist $a, b \in A$ and $\omega \in \text{Hom}(a, b)$ such that

$$(2.5) \quad |\{\alpha \in R_+^a \mid \omega(\alpha) \in -R_+^b\}| = m.$$

The latter holds for $m = 1$, since $\omega = \sigma_i^a$ fulfills Eq. (2.5) for all $a \in A$, $i \in I$, and $b = \rho_i(a)$. Suppose now that $m \in \mathbb{N}$, $a, b \in A$, and $\omega \in \text{Hom}(a, b)$ such that Eq. (2.5) holds. Since $|R^a| = \infty$, there exists $\alpha \in R_+^a$ with $\omega(\alpha) \in R_+^b$. Since ω is linear, there exists $i \in I$ such that $\omega(\alpha_i) \in R_+^b$. Then $\ell(\omega\sigma_i^{\rho_i(a)}) = m + 1$ by [HY08, Cor. 3], and hence

$$|\{\alpha \in R_+^{\rho_i(a)} \mid \omega\sigma_i^{\rho_i(a)}(\alpha) \in -R_+^b\}| = m + 1$$

by [HY08, Lemma 8(iii)]. Thus the induction step is proved.

Finally, the equivalence of (3) and (4) follows from the equivalence of (1) and (4) and the equation $\mathcal{W}(\mathcal{R}^{\text{re}}) = \mathcal{W}(\mathcal{R})$, see Prop. 2.9. $\square$

Now we prove that if $\mathcal{R}$ is a finite root system of type $\mathcal{C}$, then all roots are real, that is, $\mathcal{R}$ is uniquely determined by $\mathcal{C}$.

Proposition 2.12. *Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme and $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ a root system of type $\mathcal{C}$. Let $a \in A$, $m \in \mathbb{N}_0$, and $i_1, \dots, i_m \in I$ such that $\omega = 1_a \sigma_{i_1} \sigma_{i_2} \cdots \sigma_{i_m}$ and $\ell(\omega) = m$. Then the elements*

$$\beta_n = 1_a \sigma_{i_1} \sigma_{i_2} \cdots \sigma_{i_{n-1}}(\alpha_{i_n}) \in R_+^a,$$

where $n \in \{1, 2, \dots, m\}$ (and $\beta_1 = \alpha_{i_1}$), are pairwise different. In particular, if $\mathcal{R}$ is finite and $\omega \in \text{Hom}(\mathcal{W}(\mathcal{R}))$ is a longest element, see [HY08, Cor. 5], then

$$\{\beta_n \mid 1 \leq n \leq \ell(\omega) = |R^a|/2\} = R_+^a.$$

Proof. For all $n \in \mathbb{N}_0$ with $n \leq m$ let $a_n = \rho_{i_n} \cdots \rho_{i_2} \rho_{i_1}(a)$. Since $\ell(\sigma_{i_1}^{a_1} \sigma_{i_2}^{a_2} \cdots \sigma_{i_n}^{a_n}) = n$ for all $n \in \{1, 2, \dots, m\}$, one has $\beta_n \in R_+^a$ for these n by [HY08, Cor. 3]. By the same argument one has for all $k, n \in \mathbb{N}$ with $k < n \leq m$ the relation

$$\sigma_{i_{k+1}}^{a_{k+1}} \sigma_{i_{k+2}}^{a_{k+2}} \cdots \sigma_{i_{n-1}}^{a_{n-1}}(\alpha_{i_n}) \in R_+^{a_k}.$$

Thus $\sigma_{i_k} \sigma_{i_{k+1}} \cdots \sigma_{i_{n-1}} 1_{a_{n-1}}(\alpha_{i_n}) \neq \alpha_{i_k}$, and hence $\beta_n \neq \beta_k$ for all $k, n \in \mathbb{N}$ with $k < n \leq m$. $\square$

For any groupoid G and any $a \in \text{Ob}(G)$ let $\text{Hom}(a) = \text{Hom}(a, a) \subset \text{Hom}(G)$. Then $\text{Hom}(a)$ is a subgroup of G , which depends on a . However, the following is true.

Proposition 2.13. *Let G be a connected groupoid and $a, b \in \text{Ob}(G)$. Then $\text{Hom}(a)$ and $\text{Hom}(b)$ are isomorphic groups.*

Proof. Choose $X_b \in \text{Hom}(a, b)$. This exists since G is connected. Then the map

$$\phi_{a,b} : \text{Hom}(a) \rightarrow \text{Hom}(b), \quad g \mapsto X_b g X_b^{-1}$$

is a group homomorphism with inverse given by $\phi_{a,b}^{-1}(g) = X_b^{-1} g X_b$. $\square$

The map $\phi_{a,b}$ in the previous proof is a piece of a more general structure. Namely, let G be a connected groupoid, $a \in \text{Ob}(G)$, and for each $b \in \text{Ob}(G)$ let $X_b \in \text{Hom}(a, b)$ be a fixed morphism. Then the assignment $F_{a,X} : G \rightarrow \text{Hom}(a)$,

$$(2.6) \quad \begin{aligned} F_{a,X}(b) &= a && \text{for all } b \in \text{Ob}(G), \\ F_{a,X}(g) &= X_c^{-1} g X_b && \text{for all } g \in \text{Hom}(b, c), \end{aligned}$$

defines a fully faithful functor. In fact, G is as a groupoid isomorphic to the transformation groupoid $H := \text{Hom}(a) \times \text{Ob}(G)$ given by $\text{Ob}(H) = \text{Ob}(G)$, $\text{Hom}(H) = \text{Ob}(G) \times \text{Hom}(a) \times \text{Ob}(G)$ with composition

$$(b, g, c)(b', g', c') = \begin{cases} \text{not defined} & \text{if } c \neq b', \\ (b, gg', c') & \text{if } c = b'. \end{cases}$$

The isomorphism $G \rightarrow H$ is given by $g \mapsto (c, F_{a,X}(g), b)$ for $g \in \text{Hom}(b, c)$. In particular, G is uniquely determined by the cardinality of $\text{Ob}(G)$ and by $\text{Hom}(a)$ for any $a \in \text{Ob}(G)$.

If a connected groupoid G is presented by generators and relations, then for any $a \in \text{Ob}(G)$ the group $\text{Hom}(a)$ also can be presented by generators and relations. To do so, let $F_{a,X} : G \rightarrow \text{Hom}(a)$ be the functor defined above. The following proposition then follows from the discussion above.

Proposition 2.14. *Let G be a connected groupoid and let $a \in \text{Ob}(G)$. Suppose that J, K are index sets and $\text{Hom}(G)$ is generated by $s_j \in \text{Hom}(a_j, b_j)$, where $j \in J$, and relations $r_k = 1_{c_k} \in \text{Hom}(c_k)$, where $k \in K$. Then $\text{Hom}(a)$ is generated by $F_{a,X}(s_j)$, where $j \in J$, and relations $F_{a,X}(r_k) = 1$, where $k \in K$ and 1 is the neutral element of $\text{Hom}(a)$.*

3. STANDARD CARTAN SCHEMES AND THEIR WEYL GROUPOIDS

In this section we study root systems of type $\mathcal{C}$, where the Cartan matrices are identical for all objects. The structure of more general root systems of type $\mathcal{C}$ seems to be much more complicated, as the classification results in the next sections show.

Definition 3.1. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme such that $C^a = C^b$ for all $a, b \in A$. Then we say that $\mathcal{C}$ is *standard*, and that the Weyl groupoid $\mathcal{W}(\mathcal{C})$ is standard. If $\mathcal{R}$ is a root system of type $\mathcal{C}$, then we say that $\mathcal{R}$ is standard, if $\mathcal{C}$ is a standard Cartan scheme.

The standard Cartan schemes in the next example show that the class of root systems of type $\mathcal{C}$, where $\mathcal{C}$ is running over all Cartan schemes, is richer than the one of finite groups.

Example 3.2. Let H be a finite group. Then there exists a connected Cartan scheme $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ and a finite root system $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ of type $\mathcal{C}$, such that $\text{Hom}(a) \cong H$ for all $a \in A$. Indeed, H can be considered as a subgroup of a symmetric group $\mathbb{S}_{n+1}$. Let A be the set of left cosets gH , where $g \in \mathbb{S}_{n+1}$, and let $I = \{1, 2, \dots, n\}$. For all $gH \in A$ and $i \in I$ let $\rho_i(a) = (i, i+1)gH$, where $(i, i+1)$ is the transposition of i and $i+1$, and $C^{gH} = (c_{ij}^{gH})_{i,j \in I}$ with

$$c_{ij}^{gH} = \begin{cases} 2 & \text{if } i = j, \\ -1 & \text{if } |i - j| = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Then $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ is a standard Cartan scheme. For all $a \in A$ let $R^a = R_+^a \cup -R_+^a$, where

$$(3.1) \quad R_+^a := \{\alpha_i + \alpha_{i+1} + \dots + \alpha_j \mid 1 \leq i \leq j \leq n\},$$

be the set of roots associated to $\mathbb{S}_{n+1}$. Then $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ is a root system of type $\mathcal{C}$, and $\text{Hom}(eH) \subset \text{Hom}(\mathcal{W}(\mathcal{R}))$ is isomorphic to H .

The structure of finite connected standard root systems of type $\mathcal{C}$ is very close to the structure of reduced root systems in the sense of [Bou68, Ch. VI, §1.4].

Theorem 3.3. Let I be a non-empty finite set, $C = (c_{ij})_{i,j \in I}$ a generalized Cartan matrix, and $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ a connected standard Cartan scheme with $C^a = C$ for all $a \in A$, and let $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ be a root system of type $\mathcal{C}$.

(1) For all $a \in A$ the set $\cup_{b \in A} \text{Hom}(a, b) \subset \text{Aut}(\mathbb{Z}^\theta)$ is a group, and as such it is isomorphic to the Weyl group $W(C)$ associated to the generalized Cartan matrix C .

(2) $\mathcal{R}$ is finite if and only if C is of finite type.

(3) Assume that $\mathcal{R}$ is finite. Then for all $a \in A$, R^a is the set of roots corresponding to $W(C)$, and hence independent of the choice of $a \in A$.

Proof. Since $\mathcal{C}$ is standard, the maps $\sigma_i^a \in \text{Aut}(\mathbb{Z}^\theta)$ do not depend on the object $a \in A$, and generate the Weyl group $W(C) \subset \text{Aut}(\mathbb{Z}^\theta)$ associated to the generalized Cartan matrix C . Let $a \in A$. Since

$$\cup_{b \in A} \text{Hom}(a, b) = \{(\rho_{i_1} \cdots \rho_{i_n}(a), \sigma_{i_1} \cdots \sigma_{i_n} 1_a, a) \mid n \in \mathbb{N}_0, i_1, \dots, i_n \in I\},$$

Thm. 2.6 implies that (1) holds.

Assume that $\mathcal{R}$ is finite. Since $\mathcal{C}$ is standard, Prop. 2.12 tells that R_+^a is the set of positive roots corresponding to $W(C)$. This implies (3).

Now we prove (2). If C is of finite type, then $W(C)$ is finite. Since $\mathcal{C}$ is connected, A is finite by Part (1). Thus $\mathcal{W}(\mathcal{C})$ is finite by Part (1), and hence $\mathcal{R}$ is finite by Lemma 2.11.

Conversely, assume that $\mathcal{R}$ is finite. Then Lemma 2.11 implies that $\mathcal{W}(\mathcal{R})$ is finite, and hence $W(C)$ is finite by Part (1). Thus C is of finite type. $\square$

4. DECOMPOSITION OF FINITE ROOT SYSTEMS

In this section we study the reducibility of root systems of type $\mathcal{C}$. An analogous notion exists for root systems, see [Bou68, Ch.6, §1.2], and it is crucial for classification results.

Definition 4.1. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme. Let $J \subset I$ be a non-empty subset, and identify $\{\alpha_i \mid i \in J\}$ with the standard basis of $\mathbb{Z}^J$. For all $a \in A$ let $C'^a = (c_{ij}^a)_{i,j \in J}$. Then $\mathcal{C}' = \mathcal{C}'(J, A, (\rho_i)_{i \in J}, (C'^a)_{a \in A})$ is a Cartan scheme, called the *restriction of $\mathcal{C}$ to J* , and will be denoted by $\mathcal{C}|_J$.

Let $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ be a root system of type $\mathcal{C}$. Define $R'^a = R^a \cap \sum_{i \in J} \mathbb{Z} \alpha_i$. Then $\mathcal{R}' = \mathcal{R}'(\mathcal{C}|_J, (R'^a)_{a \in A})$ is a root system of type $\mathcal{C}|_J$, and will be denoted by $\mathcal{R}|_J$.

Restrictions are helpful to decide if a root system of type $\mathcal{C}$ is standard.

Remark 4.2. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme and $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ a root system of type $\mathcal{C}$. Assume that for each pair $(i, j) \in I \times I$ with $i \neq j$ there exists a subset $J \subset I$ such that $i, j \in J$ and $\mathcal{R}|_J$ is standard. Then $\mathcal{R}$ is standard. Indeed, $\mathcal{R}$ is standard if and only if $c_{ij}^a = c_{ij}^b$ for all $a, b \in A$ and $i, j \in I$. The latter holds by

assumption on the pairs $(i, j) \in I \times I$, and since $c_{ii}^a = 2$ for all $i \in I$ and $a \in A$.

Definition 4.3. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a Cartan scheme. Assume that $I', I'' \subset I$ are non-empty disjoint subsets such that $I = I' \cup I''$ and $c_{ij}^a = 0$ for all $i \in I', j \in I''$. Then we write $\mathcal{C} = \mathcal{C}|_{I'} \oplus \mathcal{C}|_{I''}$, and say that $\mathcal{C}$ is the *direct sum of $\mathcal{C}|_{I'}$ and $\mathcal{C}|_{I''}$* .

Let $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ be a root system of type $\mathcal{C}$. Assume that

$$R^a = \left(R^a \cap \sum_{i \in I'} \mathbb{Z} \alpha_i \right) \cup \left(R^a \cap \sum_{j \in I''} \mathbb{Z} \alpha_j \right) \quad \text{for all } a \in A.$$

Then we write $\mathcal{R} = \mathcal{R}|_{I'} \oplus \mathcal{R}|_{I''}$, and $\mathcal{R}$ is called the *direct sum of $\mathcal{R}|_{I'}$ and $\mathcal{R}|_{I''}$* . We also say that $\mathcal{R}$ is *reducible*. If $\mathcal{R} \neq \mathcal{R}|_{I_1} \oplus \mathcal{R}|_{I_2}$ for all non-empty disjoint subsets $I_1, I_2 \subset I$, then $\mathcal{R}$ is termed *irreducible*.

From now on let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a connected Cartan scheme and $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ a root system of type $\mathcal{C}$. We are going to give criteria for the reducibility of $\mathcal{R}$.

Lemma 4.4. *Let $a \in A$ and $i, j \in I$ with $i \neq j$. The following are equivalent.*

- (1) $c_{ij}^a = c_{ji}^a = 0$.
- (2) $R^a \cap (\mathbb{N}_0 \alpha_i + \mathbb{N}_0 \alpha_j) = \{\alpha_i, \alpha_j\}$.
- (3) $m_{i,j}^a = 2$.

Proof. Since $\alpha_i, \alpha_j \in R^a$ by (R2), (2) is equivalent to (3). Further, from (R1)–(R3) and Eq. (2.1) we conclude that (2) implies (1). Assume now that (1) holds, and let $\alpha := r_i \alpha_i + r_j \alpha_j \in R_+^a$, where $r_i, r_j \in \mathbb{N}_0$. Then (R1) and relation $\sigma_i^a(\alpha) = -r_i \alpha_i + r_j \alpha_j \in R^{\rho_i(a)}$ imply that $r_i = 0$ or $r_j = 0$. Hence $\alpha \in \{\alpha_i, \alpha_j\}$ by (R2). This proves (1) $\Rightarrow$ (2). $\square$

Lemma 4.5. *Suppose that $a \in A$ and $i, j \in I$, where $i \neq j$. If $c_{ij}^a = 0$ then $c_{jl}^a = c_{jl}^{\rho_i(a)}$ for all $l \in I$.*

Proof. Let $l \in I$. If $l = j$, then $c_{jl}^a = c_{jl}^{\rho_i(a)} = 2$, and if $l = i$, then $c_{jl}^a = c_{ij}^a = 0 = c_{ij}^{\rho_i(a)} = c_{jl}^{\rho_i(a)}$ by (M2) and (C2). Assume now that $l \in I \setminus \{i, j\}$. Then $\sigma_i^{\rho_j(a)} \sigma_j^a(\alpha_l) = \sigma_j^{\rho_i(a)} \sigma_i^a(\alpha_l)$ by Thm. 2.6. Explicit calculation gives

$$\sigma_i^{\rho_j(a)} \sigma_j^a(\alpha_l) = \sigma_i^{\rho_j(a)}(\alpha_l - c_{jl}^a \alpha_j) = \alpha_l - c_{il}^{\rho_j(a)} \alpha_i - c_{jl}^a \alpha_j,$$

and similarly $\sigma_j^{\rho_i(a)} \sigma_i^a(\alpha_l) = \alpha_l - c_{jl}^{\rho_i(a)} \alpha_j - c_{il}^a \alpha_i$. Comparing the coefficients of α_j gives the claim of the lemma. $\square$

Proposition 4.6. *Let $I' \subset I$ be a subset of I , and let $I'' = I \setminus I'$. Assume that $I', I'' \neq \emptyset$. The following are equivalent.*

- (1) *There exists $a \in A$ such that $c_{ij}^a = 0$ for all $i \in I'$ and $j \in I''$.*
- (2) *For all $a \in A$ and $i \in I', j \in I''$ one has $c_{ij}^a = c_{ji}^a = 0$.*
- (3) *For all $a \in A$ let $\hat{R}^a = (R^a \cap \sum_{i \in I'} \mathbb{Z}\alpha_{i'}) \cup (R^a \cap \sum_{i \in I''} \mathbb{Z}\alpha_{i''})$. Then $\hat{\mathcal{R}} = \hat{\mathcal{R}}(\mathcal{C}, (\hat{R}^a)_{a \in A})$ is a root system of type $\mathcal{C}$.*

If $\mathcal{R}$ is finite then (1)–(3) are equivalent to the following.

- (4) $\hat{\mathcal{R}} = \mathcal{R}$, where $\hat{\mathcal{R}}$ is as in (3).
- (5) $\mathcal{R} = \mathcal{R}|_{I'} \oplus \mathcal{R}|_{I''}$ with respect to the permutation $\phi = \text{id}$ of A .

Proof. The implication (2) $\Rightarrow$ (1) is trivial.

(1) $\Rightarrow$ (2). Let $b \in A$ and $l \in I'$, and assume that $c_{ij}^b = c_{ji}^b = 0$ for all $i \in I', j \in I''$. Since $c_{ij}^b = 0$ for all $j \in I''$, Lemma 4.5 implies that $c_{ji}^{\rho_l(b)} = c_{ij}^b = 0$ for all $i \in I'$ and $j \in I''$, that is, $c_{ij}^{\rho_l(b)} = c_{ji}^{\rho_l(b)} = 0$ for all $i \in I', j \in I''$. Together with the analogous argument for $l \in I''$ we obtain that $c_{ij}^{\rho_l(b)} = c_{ji}^{\rho_l(b)} = 0$ for all $i \in I', j \in I''$, and $l \in I$. Thus (2) follows from (1) since $\mathcal{R}$ is connected.

(3) $\Rightarrow$ (2). Since $\hat{R}^a \cap (\mathbb{N}_0\alpha_i + \mathbb{N}_0\alpha_j) = \{\alpha_i, \alpha_j\}$ for all $i \in I', j \in I''$, and $a \in A$, (2) follows from Lemma 4.4 and (3).

(2) $\Rightarrow$ (3). Since $\mathcal{R}$ is a root system of type $\mathcal{C}$, (2) and Lemma 4.4 imply that $m_{i,j}^a = 2$ for all $a \in A, i \in I',$ and $j \in I''$. Thus (3) is equivalent to the fact that $\sigma_l^a(\hat{R}^a) \subset \hat{R}^{\rho_l(a)}$ for all $l \in I$ and $a \in A$. Let $\alpha \in \hat{R}^a$. Then (2) implies that

$$\sigma_l^a(\alpha) \in R^{\rho_l(a)} \cap \left(\sum_{i' \in I'} \mathbb{Z}\alpha_{i'} \cup \sum_{i'' \in I''} \mathbb{Z}\alpha_{i''} \right) = \hat{R}^{\rho_l(a)},$$

and hence (3) holds.

Assume now that $\mathcal{R}$ is finite. Then, by Prop. 2.12, (2) implies that $\hat{R}_+^a = R_+^a$ for all $a \in A$, that is, (4) holds. Obviously, (4) implies (3), and (4) is also equivalent to (5), hence the proposition is proven. $\square$

For the equivalence (3) $\Leftrightarrow$ (4) in Prop. 4.6 the finiteness assumption on $\mathcal{R}$ is necessary, as the following example shows.

Example 4.7. Let $I = \{1, 2, 3, 4\}$, $A = \{a\}$, and

$$C^a = \begin{pmatrix} 2 & -2 & 0 & 0 \\ -2 & 2 & 0 & 0 \\ 0 & 0 & 2 & -2 \\ 0 & 0 & -2 & 2 \end{pmatrix},$$

$$\begin{aligned} R_+^a = & \{m\alpha_1 + (m+1)\alpha_2, (m+1)\alpha_1 + m\alpha_2, \\ & m\alpha_3 + (m+1)\alpha_4, (m+1)\alpha_3 + m\alpha_4, \mid m \in \mathbb{N}_0\} \\ & \cup \{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4\}. \end{aligned}$$

Then (2) holds in Prop. 4.6 for $I' = \{1, 2\}$ and $I'' = \{3, 4\}$. However, $R^a \neq R^a|_{\{1,2\}} \cup R^a|_{\{3,4\}}$ since $\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 \notin \hat{R}^a$, and hence $\mathcal{R}$ is irreducible.

We continue with some general statements about $\mathcal{R}$.

Lemma 4.8. *Suppose that $a \in A$ and $i, j \in I$, where $i \neq j$. The following are equivalent.*

- (1) $c_{ij}^a = c_{ji}^a = -1$.
- (2) $R^a \cap (\mathbb{N}_0\alpha_i + \mathbb{N}_0\alpha_j) = \{\alpha_i, \alpha_i + \alpha_j, \alpha_j\}$.
- (3) $m_{i,j}^a = 3$.

Proof. To (1) $\Rightarrow$ (2). Let $\alpha = c_1\alpha_i + c_2\alpha_j$ with $c_1, c_2 \in \mathbb{N}$, and assume that $\alpha \in R^a$. Then $\sigma_i^a(\alpha) = (c_2 - c_1)\alpha_i + c_2\alpha_j$, and hence relation $\sigma_i^a(\alpha) \in R_+^{\rho_i(a)} \cup -R_+^{\rho_i(a)}$ tells that $c_2 \geq c_1$. By symmetry one gets $c_1 = c_2$, and hence $\sigma_i^a(\alpha) = c_2\alpha_j$. By (R2) one obtains that $c_2 = 1$.

The implication (2) $\Rightarrow$ (3) follows from the definition of $m_{i,j}^a$. We have to prove (3) $\Rightarrow$ (1). Assume that $m_{i,j}^a = 3$. Then $c_{ij}^a, c_{ji}^a < 0$ by Lemma 4.4. Thus, $\sigma_i^{\rho_i(a)}(\alpha_j) = \alpha_j - c_{ij}^{\rho_i(a)}\alpha_i \in R_+^a \setminus \{\alpha_i, \alpha_j\}$, and hence $\beta_1 := \alpha_j - c_{ij}^a\alpha_i \in R_+^a \setminus \{\alpha_i, \alpha_j\}$ by (C2). Similarly, $\beta_2 := \alpha_i - c_{ji}^a\alpha_j \in R_+^a \setminus \{\alpha_i, \alpha_j\}$, and therefore (3) implies that $\beta_1 = \beta_2$, that is, (1) holds. $\square$

Lemma 4.9. *Suppose that $a \in A$ and $i, j \in I$ such that $m_{i,j}^a = 3$. Then $c_{il}^{\rho_i(a)} + c_{jl}^{\rho_i(a)} = c_{il}^{\rho_i\rho_j(a)} + c_{jl}^{\rho_i\rho_j(a)}$ for all $l \in I$.*

Proof. If $l \in \{i, j\}$, then Lemma 4.8 implies that both sides of the claimed equation are equal to 1. Assume now that $l \in I \setminus \{i, j\}$. Then

$$\sigma_i^{\rho_j\rho_i(a)}\sigma_j^{\rho_i(a)}\sigma_i^a(\alpha_l) = \sigma_j^{\rho_i\rho_j(a)}\sigma_i^{\rho_j(a)}\sigma_j^a(\alpha_l)$$

by Thm. 2.6, that is,

$$\begin{aligned} \alpha_l - (c_{jl}^{\rho_i(a)} + c_{il}^{\rho_j\rho_i(a)})\alpha_i - (c_{jl}^{\rho_i(a)} + c_{il}^a)\alpha_j \\ = \alpha_l - (c_{il}^{\rho_j(a)} + c_{jl}^a)\alpha_i - (c_{il}^{\rho_j(a)} + c_{jl}^{\rho_i\rho_j(a)})\alpha_j. \end{aligned}$$

One obtains the claim of the lemma by comparing the coefficients of α_j and by using (C2). $\square$

Lemma 4.10. *Suppose that $\mathcal{R}$ is finite. Let $a, b \in A$ and $i, j, l \in I$ such that $i \neq j$, $\rho_i(a) = \rho_j(a) = b \neq a$, and $\rho_l(a) = a$. Then the following hold.*

- (1) $c_{ij}^a c_{il}^a c_{jl}^a = 0$.
- (2) If $\rho_i \rho_l(b) \neq \rho_l(b)$ and $\rho_j \rho_l(b) \neq \rho_l(b)$ then $c_{ln}^a = c_{ln}^b$ for all $n \in I$.

Proof. Note that $c_{ji}^b = c_{ji}^a$ and $c_{jl}^b = c_{jl}^a$ by (C2).

Let $\tilde{\sigma} = \sigma_l^a \sigma_j^b \sigma_i^a \in \text{Hom}(a)$. Since $\mathcal{R}$ is finite, $\tilde{\sigma}$ must have finite order by Lemma 2.11. Let $W = \text{span}_{\mathbb{Z}}\{\alpha_i, \alpha_j, \alpha_l\}$. Since $\tilde{\sigma}(W) \subset W$, $\sigma := \tilde{\sigma}|_W \in \text{End}(W)$ has to have finite order as well. This will yield both Claim (1) and Claim (2).

To (1). One has $\sigma(\alpha_i) \in -\alpha_i - \mathbb{N}_0 \alpha_j - \mathbb{N}_0 \alpha_l$. If $\sigma(\alpha_i) = -\alpha_i$, then $c_{li}^a = c_{il}^a = 0$ and hence (1) holds.

It remains to consider the case when $\sigma \neq \pm \text{id}$. Then finiteness of the order of σ tells that $|\text{tr } \sigma| \in \{0, 1, 2\}$. Explicit calculation gives that

$$(4.1) \quad \text{tr } \sigma = -c_{il}^a c_{ji}^a c_{lj}^a + c_{ij}^a c_{ji}^a + c_{il}^a c_{li}^a + c_{jl}^a c_{lj}^a - 3.$$

Exchanging i and j further implies that

$$-2 \leq -c_{ij}^a c_{jl}^a c_{li}^a + c_{ij}^a c_{ji}^a + c_{il}^a c_{li}^a + c_{jl}^a c_{lj}^a - 3 \leq 2.$$

If $c_{ij}^a c_{il}^a c_{jl}^a \neq 0$, then the latter relations and (M1), (M2) imply that

$$c_{ij}^a = c_{ji}^a = c_{il}^a = c_{li}^a = c_{jl}^a = c_{lj}^a = -1.$$

In this case one has

$$\sigma = \begin{pmatrix} -1 & 1 & 1 \\ -1 & 0 & 2 \\ -2 & 1 & 2 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} -2 & 0 & 3 \\ -3 & 1 & 3 \\ -3 & 0 & 4 \end{pmatrix}$$

with respect to the basis $\{\alpha_i, \alpha_j, \alpha_l\}$. Then $\text{tr } \sigma^2 = 3$, but $\sigma^2 \neq \text{id}$, and hence σ does not have finite order. This is a contradiction to the assumption that $\mathcal{R}$ is finite, and hence (1) holds.

To (2). Assume that $\rho_i \rho_l(b) \neq \rho_l(b)$ and $\rho_j \rho_l(b) \neq \rho_l(b)$. If $c_{il}^a c_{jl}^a = 0$ then Claim (2) is valid by Lemma 4.5. Thus by (1) it remains to consider the setting $c_{ij}^a = 0$, $c_{il}^a c_{jl}^a \neq 0$. In this case one has

$$\sigma = \begin{pmatrix} -1 & 0 & -c_{il}^a \\ 0 & -1 & -c_{jl}^a \\ c_{li}^a & c_{lj}^a & t-1 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 1 - c_{il}^a c_{li}^a & -c_{il}^a c_{lj}^a & * \\ * & 1 - c_{jl}^a c_{lj}^a & * \\ * & * & t^2 - 3t + 1 \end{pmatrix}$$

with respect to the basis $\{\alpha_i, \alpha_j, \alpha_l\}$, where $t = c_{il}^a c_{li}^a + c_{jl}^a c_{lj}^a$. In the next paragraph we prove that $t \geq 4$. This implies that

$$\text{tr } \sigma^2 = t^2 - 4t + 3 = (t - 2)^2 - 1 \geq 3.$$

Since $\sigma^2 \neq \text{id}$, part (2) of the lemma is proven.

Recall that $c_{il}^a \neq 0$. Further, relation $\rho_i \rho_l(b) \neq \rho_l(b)$ implies that $\rho_i \rho_l \rho_i(a) \neq \rho_l \rho_i \rho_l(a)$, and hence $m_{i,l}^a > 3$ by Lemma 4.4 and by (R4), that is $c_{il}^a c_{li}^a \geq 2$ by Lemma 4.8. Similarly one gets $c_{jl}^a c_{lj}^a \geq 2$. Thus the assumptions in part (2) imply that $t \geq 4$. $\square$

Now we present another technique for the analysis of the finiteness of $\mathcal{W}(\mathcal{R})$. We will use this method in Sect. 6.

Proposition 4.11. *Let $m, n \in \mathbb{N}$, and define the families $G_{m,n}$ and $H_{m,n}$ of groups by generators and relations as follows.*

$$\begin{aligned} G_{m,n} &= \langle s, t \rangle / (s^2, t^m, (st^{-1}st)^n), \\ H_{m,n} &= \langle s_1, \dots, s_m, T \rangle / (s_i^2, T^m, (s_i s_{i+1})^n, T^{-1} s_i T s_{i+1} \mid 1 \leq i \leq m), \end{aligned}$$

where the convention $s_{m+1} = s_1$ is used in the definition of $H_{m,n}$. Then there is a group isomorphism $\varphi : G_{m,n} \rightarrow H_{m,n}$ with $\varphi(s) = s_m$, $\varphi(t) = T$. Further, $G_{m,n}$ is finite if and only if $m = 1$ or $m = 2$ or $n = 1$ or $(m, n) = (3, 2)$.

Proof. Since $T^{-1} s_m T = s_1$ and $(s_m s_1)^n = 1$ in $H_{m,n}$, there is a unique group homomorphism $\varphi : G_{m,n} \rightarrow H_{m,n}$ with $\varphi(s) = s_m$, $\varphi(t) = T$. Further, there is a group homomorphism $\psi : H_{m,n} \rightarrow G_{m,n}$ with $\psi(s_i) = t^{-i} s t^i$, $\psi(T) = t$, and the identities $\varphi\psi = \text{id}$ and $\psi\varphi = \text{id}$ hold. Thus φ is an isomorphism.

If $m = 1$ then $H_{m,n} \simeq \mathbb{Z}/2\mathbb{Z}$. Assume now that $m \geq 2$. Let $N := \langle s_i \mid 1 \leq i \leq m \rangle \subset H_{m,n}$. Since $T^{-1} s_i T = s_{i+1}$ for all $i \in \{1, 2, \dots, m\}$, N is a normal subgroup of $H_{m,n}$, and $H_{m,n}$ is the semidirect product of N and the finite abelian group $\langle T \rangle / (T^m) \simeq \mathbb{Z}/m\mathbb{Z}$. Thus $H_{m,n}$ is finite if and only if N is finite. But

$$N = \langle s_1, \dots, s_m \rangle / (s_i^2, (s_i s_{i+1})^n \mid 1 \leq i \leq m)$$

is a Coxeter group. It is easy to see that N is finite if and only if $n = 1$ (and $N \simeq \mathbb{Z}/2\mathbb{Z}$) or $m = 2$ (and $N \simeq \mathbb{D}_n$, the dihedral group of order $2n$) or $(m, n) = (3, 2)$ (and $N \simeq (\mathbb{Z}/2\mathbb{Z})^3$). $\square$

Proposition 4.12. *Let $m, n, p \in \mathbb{N}$, and define the families $G_{m,n,p}$ and $H_{m,n,p}$ of groups by generators and relations as follows.*

$$\begin{aligned} G_{m,n,p} &= \langle s, u, t \rangle / (s^2, u^2, t^m, (st^{-1}ut)^n, (su)^p), \\ H_{m,n,p} &= \langle s_1, \dots, s_m, u_1, \dots, u_m, T \rangle / (s_i^2, u_i^2, T^m, (s_i u_{i+1})^n, (s_i u_i)^p, \\ &\quad T^{-1} s_i T s_{i+1}, T^{-1} u_i T u_{i+1} \mid 1 \leq i \leq m), \end{aligned}$$

where the convention $s_{m+1} = s_1$, $u_{m+1} = u_1$ is used in the definition of $H_{m,n,p}$. Then there is a group isomorphism $\varphi : G_{m,n,p} \rightarrow H_{m,n,p}$ with $\varphi(s) = s_m$, $\varphi(u) = u_m$, $\varphi(t) = T$. Further, $G_{m,n,p}$ is finite if and only if $m = 1$ or $(n, p) = (1, 1)$ or $(m, n) = (2, 1)$ or $(m, p) = (2, 1)$ or $(m, n, p) = (3, 1, 2)$ or $(m, n, p) = (3, 2, 1)$.

Proof. Entirely similar to the proof of Prop. 4.11. $\square$

5. ROOT SYSTEMS OF TYPE $\mathcal{C}$ WITH TWO OBJECTS

Let $I = \{1, 2, \dots, \theta\}$ for some $\theta \in \mathbb{N}$, and $A = \{x, y\}$ with $x \neq y$. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a connected Cartan scheme and $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ a root system of type $\mathcal{C}$. Without loss of generality suppose that

$$(5.1) \quad \begin{aligned} \rho_i(x) &= y, & \rho_i(y) &= x & \text{if } 1 \leq i \leq \kappa, \\ \rho_i(x) &= x, & \rho_i(y) &= y & \text{if } \kappa + 1 \leq i \leq \theta \end{aligned}$$

for some $\kappa \in I$. In this case (C2) implies that $c_{ij}^x = c_{ij}^y$ whenever $1 \leq i \leq \kappa$ and $j \in I$.

If $\theta = 1$ then $\rho_1(x) = y$, $R^x = \{\alpha_1, -\alpha_1\}$, $R^y = \{\alpha_1, -\alpha_1\}$, and $C^x = C^y = (2)$, see also [HY08, Ex. 1].

Consider now the case $\theta = 2$ and $\kappa = 1$. Then $\mathcal{W}(\mathcal{R})$ is isomorphic to the Coxeter groupoid

$$\langle s_1^x, s_1^y, s_2^x, s_2^y \rangle / (s_1^y s_1^x, s_1^x s_1^y, (s_2^x)^2, (s_2^y)^2, (s_1^y s_2^y s_1^x s_2^x)^m)$$

for some $m \in \mathbb{N}$, see Thm. 2.6. Identify σ_i^a , where $i \in \{1, 2\}$ and $a \in \{x, y\}$, with its matrix with respect to the standard basis $\{\alpha_1, \alpha_2\}$ of $\mathbb{Z}^2$. Further, note that $c_{12}^x = c_{12}^y$ by (C2). One gets

$$\begin{aligned} \sigma_1 \sigma_2 \sigma_1 \sigma_2^x &= \begin{pmatrix} -1 & -c_{12}^y \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -c_{21}^y & -1 \end{pmatrix} \begin{pmatrix} -1 & -c_{12}^x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -c_{21}^x & -1 \end{pmatrix} \\ &= \begin{pmatrix} c_{12}^x c_{21}^y - 1 & c_{12}^x \\ -c_{21}^y & -1 \end{pmatrix} \begin{pmatrix} c_{12}^x c_{21}^x - 1 & c_{12}^x \\ -c_{21}^x & -1 \end{pmatrix} \\ &= \begin{pmatrix} c_{12}^x c_{21}^x c_{12}^x c_{21}^y - c_{12}^x c_{21}^y - 2c_{12}^x c_{21}^x + 1 & c_{12}^x (c_{12}^x c_{21}^y - 2) \\ -c_{12}^x c_{21}^x c_{21}^y + c_{21}^y + c_{21}^x & 1 - c_{12}^x c_{21}^y \end{pmatrix}. \end{aligned}$$

The groupoid $\mathcal{W}(\mathcal{R})$ is finite if and only if $\sigma := \sigma_1\sigma_2\sigma_1\sigma_2^x$ has finite order. Since $\sigma \in \text{Aut}(\mathbb{Z}^2)$, the latter is equivalent to the condition

$$(5.2) \quad \sigma = \pm \text{id} \quad \text{or} \quad \text{tr } \sigma \in \{-1, 0, 1\}.$$

By (M2) and (C2) one has $\sigma = \text{id}$ if and only if $c_{12}^x = c_{21}^x = c_{12}^y = c_{21}^y = 0$. Further, $\sigma = -\text{id}$ if and only if $c_{12}^x c_{21}^y = 2$, $c_{21}^x = c_{21}^y$, that is

$$C^x = C^y = \begin{pmatrix} 2 & -1 \\ -2 & 2 \end{pmatrix} \quad \text{or} \quad C^x = C^y = \begin{pmatrix} 2 & -2 \\ -1 & 2 \end{pmatrix}.$$

It remains to consider the second relation in Eq. (5.2). One has

$$\text{tr } \sigma = (c_{12}^x c_{21}^x - 2)(c_{12}^x c_{21}^y - 2) - 2,$$

and hence relation $-1 \leq \text{tr } \sigma \leq 1$ is equivalent to

$$1 \leq (c_{12}^x c_{21}^x - 2)(c_{12}^x c_{21}^y - 2) \leq 3.$$

By (M1) and (M2) one gets $c_{12}^x = c_{21}^x = c_{12}^y = c_{21}^y = -1$ or $c_{12}^x c_{21}^x \geq 3$, $c_{12}^x c_{21}^y \geq 3$. However, the first case contradicts (R4) and Lemma 4.8. Now the following statement can be obtained easily.

Proposition 5.1. *Assume that $\theta = 2$ and $\kappa = 1$. Then $\mathcal{R}$ is finite if and only if the Cartan matrices C^x, C^y satisfy, up to permutation of x and y , one of the following conditions (1),(2).*

- (1) $C^x = C^y$ is of finite type $A_1 \times A_1, B_2$, or G_2 , that is, $c_{12}^x = c_{21}^x = 0$ or $c_{12}^x c_{21}^x \in \{2, 3\}$.
- (2) $c_{12}^x = c_{12}^y = -1$, $c_{21}^x = -3$, and $c_{21}^y \in \{-4, -5\}$.

In case (1) $R^x = R^y$ is the usual set of roots corresponding to the generalized Cartan matrix $C^x = C^y$. In case (2) one has

$$R_+^x = \{1, 2, 12, 12^2, 12^3, 1^2 2^3, 1^3 2^4, 1^3 2^5\},$$

$$R_+^y = \{1, 2, 12, 12^2, 12^3, 12^4, 1^2 2^3, 1^2 2^5\}$$

if $c_{21}^y = -4$, and

$$R_+^x = \{1, 2, 12, 12^2, 12^3, 1^2 2^3, 1^3 2^4, 1^3 2^5, 1^4 2^5, 1^4 2^7, 1^5 2^7, 1^5 2^8\},$$

$$R_+^y = \{1, 2, 12, 12^2, 12^3, 12^4, 12^5, 1^2 2^3, 1^2 2^5, 1^2 2^7, 1^3 2^7, 1^3 2^8\}$$

if $c_{21}^y = -5$.

In the last part of the proposition the abbreviation $1^m 2^n = m\alpha_1 + n\alpha_2$ was used, where exponents 1 and factors i^0 for $i \in \{1, 2\}$ are omitted. The determination of R_+^a is straightforward, see Prop. 2.12 or, more directly, [HY08, Lemma 6].

The case $\theta = 2, \kappa = 2$ is even easier than the case $\kappa = 1$. Indeed, by (C2) one has $C^x = C^y$, and hence Prop. 2.12 implies that $R^x = R^y$

is the root system of rank 2 corresponding to the Cartan matrix C^x . Thus the following holds.

Proposition 5.2. *Assume that $\theta = 2$ and $\kappa = 2$. Then $\mathcal{R}$ is finite if and only if $\mathcal{R}$ is standard and $C^x = C^y$ is of finite type, that is, $c_{12}^x = c_{21}^x = 0$ or $c_{12}^x c_{21}^x \in \{1, 2, 3\}$.*

Now assume that $\theta \geq 3$. As before, suppose that $1 \leq \kappa \leq \theta$ such that Eq. (5.1) holds. First we develop some general properties.

Lemma 5.3. *Suppose that $\mathcal{R}$ is finite and that $1 \leq \kappa \leq \theta - 2$. Then $c_{li}^x c_{lj}^x = 0$ for all $l \leq \kappa$ and $i, j > \kappa$ with $i \neq j$.*

Proof. Let $I' = \{l, i, j\}$, and consider the restriction $\mathcal{R}|_{I'}$ of $\mathcal{R}$, see Sect. 4. Since $\rho_l(x) = y$, $\mathcal{R}|_{I'}$ is connected. Thus it suffices to consider the case $I = I'$. Let $t_i := \sigma_l^y \sigma_i^y \sigma_l^x$, $t_j := \sigma_l^y \sigma_j^y \sigma_l^x$. By Prop. 2.14, the subgroup $\text{Hom}(x)$ of $\text{Hom}(\mathcal{W}(\mathcal{R}))$ is generated by $s_i := \sigma_i^x$, $s_j := \sigma_j^x$, t_i , and t_j . Moreover, $\text{Hom}(x)$ can be presented by the relations

$$s_i^2, s_j^2, t_i^2, t_j^2, (s_i s_j)^{m_{i,j}^x}, (t_i t_j)^{m_{i,j}^y}, (s_i t_i)^{m_{i,i}^x/2}, (s_j t_j)^{m_{j,j}^x/2}.$$

Assume now that $c_{li}^x c_{lj}^x \neq 0$. Then $m_{l,i}^x > 2$ by Lemma 4.4, that is $m_{l,i}^x/2 \geq 2$, and similarly $m_{l,j}^x/2 \geq 2$. Thus the subgroup of $\text{Hom}(x)$ generated by s_i and t_j is infinite and hence $\text{Hom}(x)$ is an infinite Coxeter group. This is a contradiction to the finiteness of $\mathcal{R}$, and hence the claim of the lemma is proven. $\square$

Theorem 5.4. *Let $\theta \in \mathbb{N}$, $I = \{1, 2, \dots, \theta\}$, $A = \{x, y\}$, and $\kappa \in I$ as in Eq. (5.1). If $\mathcal{R}$ is finite and irreducible, then up to permutation of $\{x, y\}$ and I , one of the following sets of conditions hold.*

- (1) $\theta \in \mathbb{N}$, $\kappa \in \{1, 2, \dots, \theta\}$, and $C^x = C^y$ is an indecomposable Cartan matrix of finite type such that $c_{ij}^x c_{ji}^x \neq 1$ for all i, j with $i \leq \kappa$, $j > \kappa$.
- (2) $\theta = 2$, $\kappa = 1$, $c_{12}^x = c_{12}^y = -1$, $c_{21}^x = -3$, $c_{21}^y \in \{-4, -5\}$.
- (3) $\theta = 3$, $\kappa = 1$,

$$C^x = \begin{pmatrix} 2 & -1 & 0 \\ -2 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}, \quad C^y = \begin{pmatrix} 2 & -1 & 0 \\ -2 & 2 & -2 \\ 0 & -1 & 2 \end{pmatrix}.$$

- (4) $\theta = 3$, $\kappa = 1$,

$$C^x = \begin{pmatrix} 2 & -2 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}, \quad C^y = \begin{pmatrix} 2 & -2 & 0 \\ -1 & 2 & -2 \\ 0 & -1 & 2 \end{pmatrix}.$$

Conversely, if C^x, C^y satisfy one of the conditions (1)–(4), then $\mathcal{R}$ is finite and irreducible.

Proof. If $\theta \leq 2$ then the claim of the theorem holds by Props. 5.1, 5.2.

Assume that $\theta \geq 3$. Consider first the case $\kappa = 1$, that is, $\rho_1(x) = y$, $\rho_1(y) = x$, and $\rho_i(x) = x$, $\rho_i(y) = y$ if $2 \leq i \leq \theta$. Using a permutation of $\{2, 3, \dots, \theta\}$, by Lemma 5.3 one can assume that $c_{1i}^x = 0$ for all $i \geq 3$. Then

$$(5.3) \quad c_{il}^x = c_{il}^y \quad \text{for all } i \geq 3 \text{ and } l \in I$$

by Lemma 4.5, and hence $c_{i1}^x = c_{1i}^y = c_{i1}^y = 0$ for all $i \geq 3$ by (M2). We obtain that $\sigma_1^y \sigma_i^y \sigma_1^x = \sigma_i^x$ for all $i \geq 3$ by Thm. 2.6. Therefore Prop. 2.14 tells that $\text{Hom}(x)$ is isomorphic to the group

$$(5.4) \quad \langle s_2, s_3, \dots, s_\theta, t \rangle / (s_i^2, t^2, (s_j s_k)^{m_{j,k}^x}, (ts_l)^{m_{2,l}^y}, (s_2 t)^{m_{1,2}^x/2} \mid 2 \leq i \leq \theta, 2 \leq j < k \leq \theta, 2 < l \leq \theta),$$

where t corresponds to the element $\sigma_1^y \sigma_2^y \sigma_1^x$. Since $\mathcal{R}$ is irreducible and $m_{1,i}^x = 2$ for $i \geq 3$, we obtain that $m_{1,2}^x \geq 3$ by Lemma 4.4 and Prop. 4.6. Thus $\text{Hom}(x)$ is a Coxeter group of rank θ . Further, again since $\mathcal{R}$ is irreducible and $c_{1i}^x = 0$ for all $i \geq 3$, we have $c_{2r}^x \neq 0$ for some $r \geq 3$. Without loss of generality assume that $c_{23}^x \neq 0$. Then $c_{32}^y = c_{32}^x \neq 0$ by Eq. (5.3) and (M2), and hence $c_{23}^y \neq 0$ and $m_{2,3}^x, m_{2,3}^y \geq 3$. Since $\text{Hom}(x)$ is a finite Coxeter group, this implies that $s_2 t = t s_2$, that is, $m_{1,2}^x = 4$. Then Prop. 5.1 gives that $c_{ij}^x = c_{ij}^y$ for $i, j \in \{1, 2\}$, and $c_{12}^x c_{21}^x = 2$.

First assume that $\theta \geq 4$ and $c_{2i}^x \neq 0$ for some $i \geq 4$. As above, we conclude that $m_{2,i}^x, m_{2,i}^y \geq 3$, which is a contradiction to the finiteness of $\text{Hom}(x)$ and the relations $m_{2,3}^x, m_{2,3}^y \geq 3$. Hence, if $\theta \geq 4$, then $c_{2i}^x = c_{2i}^y = 0$ for all $i \geq 4$. Since $\mathcal{R}$ is irreducible, there exists $r \geq 4$ with $c_{3r}^x \neq 0$. Then $m_{2,3}^x = m_{2,3}^y = 3$ by the finiteness of the group in Eq. (5.4), and hence $c_{2i}^x = c_{2i}^y$ for all i . Thus, if $\theta \geq 4$, then $C^x = C^y$ as in (1). The restriction $c_{1j}^x c_{j1}^x \neq 1$ comes from Lemma 4.8 and the relation $\rho_1 \rho_j \rho_1(x) = x \neq y = \rho_j \rho_1 \rho_j(x)$ for $j > 1$.

If $\theta = 3$ and $\kappa = 1$, then Eq. (5.4) tells that

$$(5.5) \quad \text{Hom}(x) \simeq \langle s_2, s_3, t \rangle / (s_2^2, s_3^2, t^2, (s_2 s_3)^{m_{2,3}^x}, (ts_3)^{m_{2,3}^y}, s_2 t s_2 t).$$

Further, $m_{2,3}^x, m_{2,3}^y \geq 3$ by the above arguments. Thus, $(m_{2,3}^x, m_{2,3}^y) \in \{(3, 3), (3, 4), (4, 3)\}$. Since $\rho_2(a) = \rho_3(a) = a$ for $a \in \{x, y\}$, Thm. 3.3 yields that $(c_{23}^x c_{32}^x, c_{23}^y c_{32}^y) \in \{(1, 1), (1, 2), (2, 1)\}$. Recall that $c_{ij}^x = c_{ij}^y$ for all $(i, j) \neq (2, 3)$. Thus, if $c_{23}^a c_{32}^a = 1$ for $a \in \{x, y\}$, then $\mathcal{R}$ satisfies the conditions in (1). Otherwise, since $c_{32}^x = c_{32}^y$, $\mathcal{R}$ satisfies up to permutation of x and y the conditions of (3) or (4).

Consider now the case $\theta \geq 3$, $\kappa \geq 2$. From (C2) we obtain that $c_{in}^x = c_{in}^y$ for all $i \leq \kappa$ and $n \in I$. Further, Lemma 4.10(2) for $l > \kappa$, $i, j \leq \kappa$,

$a := x$, and $b := y$ implies that $c_{ln}^x = c_{ln}^y$ for all $l > \kappa$ and $n \in I$. Thus, $\mathcal{R}$ satisfies the conditions in (1), where the restriction $c_{ij}^x c_{ji}^x \neq 1$ comes again from Lemma 4.8 and the relation $\rho_i \rho_j \rho_i(x) = x \neq y = \rho_j \rho_i \rho_j(x)$ for $i \leq \kappa, j > \kappa$. $\square$

Remark 5.5. The appearance of non-standard root systems in Thm. 5.4 is not surprising. With an appropriate definition of the Weyl groupoid of a Nichols algebra of diagonal type, the examples in Thm. 5.4(2) can be identified with the root systems of the Nichols algebras corresponding to Row 14 and Row 17 of [Hec05, Table 1], respectively. Similarly, the examples in Thm. 5.4(3),(4) can be identified with the root systems of the Nichols algebras corresponding to Row 13 and Row 18 of [Hec05, Table 2], respectively.

6. ROOT SYSTEMS OF TYPE $\mathcal{C}$ WITH THREE OBJECTS

Let $\theta \in \mathbb{N}$, $I = \{1, 2, \dots, \theta\}$, and A a set of cardinality 3. Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a connected Cartan scheme and $\mathcal{R} = \mathcal{R}(\mathcal{C}, (R^a)_{a \in A})$ a root system of type $\mathcal{C}$. In this case we necessarily have $\theta \geq 2$.

Let first $\theta = 2$. Then, up to enumeration of the objects and up to permutation of I , we may fix the three elements x, y , and z of A , such that the object change diagram of $\mathcal{R}$ is

$$(6.1) \quad \begin{array}{ccccc} x & & 1 & & y & & 2 & & z \\ \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet \end{array}$$

By (C2) the generalized Cartan matrices are

$$(6.2) \quad C^x = \begin{pmatrix} 2 & -a \\ -c & 2 \end{pmatrix}, \quad C^y = \begin{pmatrix} 2 & -a \\ -d & 2 \end{pmatrix}, \quad C^z = \begin{pmatrix} 2 & -b \\ -d & 2 \end{pmatrix},$$

where $a, b, c, d \in \mathbb{N}_0$. Further — by replacing $(1, 2)$ by $(2, 1)$ and (x, z) by (z, x) , if necessary — one may assume that $a \leq d$.

Theorem 6.1. *Let $\mathcal{R}$ be a connected root system of type $\mathcal{C}$ of rank 2 with 3 objects. Assume that $A = \{x, y, z\}$ and $I = \{1, 2\}$ such that the object change diagram of $\mathcal{R}$ is as in Eq. (6.1). Further, let $a, b, c, d \in \mathbb{N}_0$ such that $a \leq d$ and that the Cartan matrices C^x, C^y and C^z are as in Eq. (6.2). If $\mathcal{R}$ is finite, then (a, b, c, d) and R^x satisfy one of the following equations.*

- (1) $(a, b, c, d) = (1, 1, 1, 1)$, $|R_+^x| = 3$.
- (2) $(a, b, c, d) = (1, 1, 3, 3)$, $|R_+^x| = 6$.
- (3) $(a, b, c, d) = (1, 2, 4, 2)$, $|R_+^x| = 6$.
- (4) $(a, b, c, d) = (1, 3, 6, 2)$, $|R_+^x| = 12$.

- (5) $(a, b, c, d) = (1, 4, 5, 2)$, $|R_+^x| = 12$.
- (6) $(a, b, c, d) = (1, 3, 7, 2)$, $|R_+^x| = 18$.
- (7) $(a, b, c, d) = (1, 5, 5, 2)$, $|R_+^x| = 18$.

Conversely, if (a, b, c, d) is one of the above 7 quadruples, then $\mathcal{R}$ is finite.

Proof. If one of the relations $a = 0$, $b = 0$, $c = 0$, and $d = 0$ holds, then also the other three because of (M2). However, then $x = \rho_2 \rho_1(y) = \rho_1 \rho_2(y) = z$ by (R4) and Lemma 4.4, which is a contradiction. Thus $a, b, c, d > 0$.

Clearly, the finiteness of $\mathcal{R}$ implies that the order of the linear map $\tilde{\sigma} := \sigma_2^x \sigma_1^y \sigma_2^z \sigma_1^z \sigma_2^y \sigma_1^x \in \text{Aut}(\mathbb{Z}^2)$ is finite, that is, the matrix $t = (t_{ij})_{i,j=1,2}$, where

$$\begin{aligned} t_{11} &= -abd^2 + 2ad + bd - 1, \\ t_{12} &= a^2bd^2 - 2a^2d - 2abd + 2a + b, \\ t_{21} &= -abcd^2 + 2acd + bcd + bd^2 - c - 2d, \\ t_{22} &= a^2bcd^2 - 2a^2cd - 2abcd - abd^2 + 2ac + 2ad + bc + bd - 1, \end{aligned}$$

has finite order. This means that either $t = \text{id}$ or $t = -\text{id}$ or $t_{11} + t_{22} \in \{-1, 0, 1\}$. Observe that

$$(6.3) \quad t_{11} + t_{22} - 2 = (ad - 1)(abcd - 2ac - bc - 2bd + 4),$$

$$(6.4) \quad t_{11} + t_{22} + 2 = (abd - 2a - b)(acd - c - 2d),$$

$$(6.5) \quad t_{22} = (1 - ac)t_{11} + c(-abd + a + b).$$

Thus, if $t = \text{id}$, then $1 - ac - abcd + ac + bc = 1$ by Eq. (6.5), that is, $ad = 1$. Therefore $a = d = 1$, and relation $t_{21} = 0$ gives that $b + c = 2$. Since $bc \neq 0$, we obtain that $b = c = 1$. This gives solution (1).

Suppose now that $t = -\text{id}$. Then $-1 + 2ac - abcd + bc = -1$ by Eq. (6.5), that is, $2a + b = abd$. Inserting this into Eq. (6.3) we get that $-4 = (ad - 1)(4 - 2bd)$. Since $d \geq a$, we obtain the solutions $(a, b, d) \in \{(1, 2, 2), (1, 1, 3)\}$. Using equation $t_{21} = 0$ we get solutions (2) and (3).

It remains to determine all solutions with $t_{11} + t_{22} \in \{-1, 0, 1\}$. Eq. (6.3) yields that $ad \in \{2, 3, 4\}$. If $ad = 4$, then either $(a, d) = (1, 4)$ or $(a, d) = (2, 2)$. Further, by Eq. (6.4) we get that

$$(3b - 2a)(3c - 2d) \in \{1, 2, 3\}.$$

Both for $(a, d) = (1, 4)$ and for $(a, d) = (2, 2)$ there is a unique solution of this relation with $b, c \in \mathbb{N}$, namely, $(a, b, c, d) = (1, 1, 3, 4)$ and $(a, b, c, d) = (2, 1, 1, 2)$. However, in the first case one has

$$(\sigma_1 \sigma_2)^4 \sigma_1^x(\alpha_1) = \alpha_1 - \alpha_2,$$

and in the second case one gets $\sigma_2\sigma_1\sigma_2\sigma_1^x(\alpha_2) = \alpha_1 - \alpha_2$. Therefore (R1) gives that there is no root system of type $\mathcal{C}$ with $ad = 4$.

Assume now that $ad = 3$, that is, $a = 1$ and $d = 3$. Then Eq. (6.3) implies that $-3 \leq 2(2bc - 2c - 6b + 4) \leq -1$, which has no solution with $a, b, c, d \in \mathbb{N}$.

Finally, let $a = 1$ and $d = 2$. Then, by Eq. (6.4), $t_{11} + t_{22} \in \{-1, 0, 1\}$ if and only if

$$(6.6) \quad 1 \leq (b - 2)(c - 4) \leq 3.$$

If $b = 1$ then $c \in \{1, 2, 3\}$. If $(b, c) = (1, 1)$ then $\sigma_2^x\sigma_1^y\sigma_2^z(\alpha_1) = \alpha_1 - \alpha_2$, a contradiction to (R1). If $(b, c) = (1, 2)$ then $C^x = C^y = C^z$, $|R_+^x| = 4$, but $(\rho_1\rho_2)^4(x) = y \neq x$, a contradiction to (R4). If $(b, c) = (1, 3)$ then $\sigma_1^z\sigma_2^y\sigma_1^x\sigma_2^x(\alpha_1) = \alpha_2 - \alpha_1$, a contradiction to (R1).

If $b = 3$ in Rel. (6.6), then $c \in \{5, 6, 7\}$. If $c = 5$ then direct computation shows that $(\sigma_1\sigma_2)^4 1_y(\alpha_1) = \alpha_2 - \alpha_1$, a contradiction to (R1). If $c = 6$, then we get solution (4), and if $c = 7$, then we get solution (6).

The remaining two solutions of Rel. (6.6) are $(b, c) = (4, 5)$ and $(b, c) = (5, 5)$. These correspond to solution (5) and (7), respectively.

The sets R_+^x can be calculated from Prop. 2.12. $\square$

Remark 6.2. It is interesting to note that in contrast to the case with two objects, see Rem. 5.5, not all non-standard root systems can be obtained from Nichols algebras of diagonal type. The example in Thm. 6.1(3) can be identified with the root system of the Nichols algebras corresponding to Row 10 [Hec05, Table 1]. However, the only rank two Nichols algebras of diagonal type with 12 positive roots are those in Row 17 of [Hec05, Table 1], and there are no such Nichols algebras with more than 12 positive roots. This can be read off from the trees in the appendix of [Hec07]. The Nichols algebras corresponding to Row 17 of [Hec05, Table 1] have been discussed already in Rem. 5.5: in all of the Cartan matrices one has at least one entry -1 . Thus the examples in Thm. 6.1(4)–(7) can not be obtained as the root system of a Nichols algebra of diagonal type.

It is not clear if there are more general Nichols algebras with such root systems.

Now we assume that $\theta = 3$.

Theorem 6.3. *Let $\mathcal{C}$ be a connected Cartan scheme with $I = \{1, 2, 3\}$ and with 3 objects, and let $\mathcal{R}$ be a finite irreducible root system of type $\mathcal{C}$. Then $\mathcal{R}$ is standard, and the Cartan matrices are indecomposable and of type A_3 , B_3 , or C_3 . If $c_{12}^a = c_{21}^a = -1$ and $c_{13}^a = c_{31}^a = 0$ for all*

objects a , then the object change diagram of $\mathcal{R}$ is

$$\begin{array}{ll} \bullet \xrightarrow{1,3} \bullet \xrightarrow{2} \bullet & \text{for type } A_3 \text{ Cartan matrices, and} \\ \bullet \xrightarrow{1} \bullet \xrightarrow{2} \bullet & \text{for type } B_3 \text{ and } C_3 \text{ Cartan matrices.} \end{array}$$

Proof. Let x, y, z denote the three objects of $\mathcal{R}$. Since $\mathcal{C}$ is connected, we may assume (using permutations of I and A) that the restriction of $\mathcal{R}$ to $I = \{1, 2\}$ is as for $\theta = 2$ — without supposing that $a \leq d$ in Eq. (6.2). Then we have to consider three cases.

Case 1: Assume that $\rho_3 : A \rightarrow A$ is the identity. By Thm. 2.6 and Prop. 2.14 the group $\text{Hom}(x)$ is isomorphic to

$$(6.7) \quad \text{Hom}(x) \simeq \langle s_2, s_3, t_3, u_1, u_3 \rangle / (s_2^2, s_3^2, t_3^2, u_1^2, u_3^2, (u_1 s_2)^{m_{1,2}^x/3}, (t_3 s_3)^{m_{1,3}^x/2}, (s_2 s_3)^{m_{2,3}^x}, (u_3 t_3)^{m_{2,3}^y/2}, (u_1 u_3)^{m_{1,3}^z}),$$

where the isomorphism is given by $\sigma_2^x \mapsto s_2$, $\sigma_3^x \mapsto s_3$, $\sigma_1^y \sigma_3^y \sigma_1^x \mapsto t_3$, $\sigma_1^y \sigma_2^z \sigma_1^z \sigma_2^y \sigma_1^x \mapsto u_1$, and $\sigma_1^y \sigma_2^z \sigma_3^z \sigma_2^y \sigma_1^x \mapsto u_3$. Thus, $\text{Hom}(x)$ is a Coxeter group.

Suppose first that $m_{1,3}^x = m_{2,3}^y = 2$. Then $c_{13}^x = c_{23}^y = 0$, and hence $c_{13}^y = 0$ by (C2) and relation $\rho_1(x) = y$. Thus, $\mathcal{R}$ is not irreducible by Prop. 4.6, which is a contradiction. Note that $m_{2,3}^z = m_{2,3}^y$, since $\rho_2(y) = z$. Now, using a symmetry in the presentation of $\mathcal{R}$, we can assume that $m_{2,3}^y > 2$.

If $m_{1,2}^x > 3$, then the quotient

$$\text{Hom}(x)/(t_3 s_3) \simeq \langle s_2, s_3, u_1, u_3 \rangle / (s_2^2, s_3^2, u_1^2, u_3^2, (u_1 s_2)^{m_{1,2}^x/3}, (s_2 s_3)^{m_{2,3}^x}, (u_3 s_3)^{m_{2,3}^y/2}, (u_1 u_3)^{m_{1,3}^z})$$

is a Coxeter group without relation between u_1 and s_3 , and hence it is infinite, which is a contradiction to the finiteness of $\mathcal{R}$. If $m_{1,2}^x = 3$, then

$$\text{Hom}(x) \simeq \langle s_2, s_3, t_3, u_3 \rangle / (s_2^2, s_3^2, t_3^2, u_3^2, (t_3 s_3)^{m_{1,3}^x/2}, (s_2 s_3)^{m_{2,3}^x}, (u_3 t_3)^{m_{2,3}^y/2}, (s_2 u_3)^{m_{1,3}^z}).$$

In this case, if $m_{1,3}^x > 2$, then there is no Coxeter relation between s_2 and t_3 , which is again a contradiction. Thus $m_{1,3}^x = 2$ and

$$(6.8) \quad \text{Hom}(x) \simeq \langle s_2, s_3, u_3 \rangle / (s_2^2, s_3^2, u_3^2, (s_2 s_3)^{m_{2,3}^x}, (u_3 s_3)^{m_{2,3}^y/2}, (s_2 u_3)^{m_{1,3}^z}).$$

Since $m_{1,2}^x = 3$, $m_{1,3}^x = 2$ and $m_{2,3}^y > 2$, Thm. 6.1 yields that the Cartan matrices C^x , C^y , and C^z are

$$C^x = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -a \\ 0 & -b & 2 \end{pmatrix}, C^y = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -c \\ 0 & -d & 2 \end{pmatrix}, C^z = \begin{pmatrix} 2 & -1 & -e \\ -1 & 2 & -c \\ -f & -g & 2 \end{pmatrix},$$

where $a, b, e, f, g \in \mathbb{N}_0$ and $c, d \in \mathbb{N}$. Moreover, Lemmata 4.5, 4.9 imply that

$$(6.9) \quad d = b, \quad a = c + e.$$

The isomorphism in Eq. (6.8) tells that σ_2^x and σ_3^x generate a finite Coxeter subgroup of $\text{Hom}(x)$, and hence $ab \in \{0, 1, 2, 3\}$. Since $a = c + e$ and $c > 0$, we get $ab \in \{1, 2, 3\}$, and hence $m_{2,3}^x \geq 3$.

If $ab = 3$, then $m_{2,3}^x = 6$, and therefore the finiteness of $\text{Hom}(x)$ and Eq. (6.8) imply that $m_{2,3}^y = 4$, $m_{1,3}^z = 2$. The former gives by Prop. 5.1 that $cd = 2$, and the latter, that $e = f = 0$. By Eq. (6.9) we get $2 = cd = ab = 3$, a contradiction.

If $ab = 1$, then $a = b = 1$, and hence Eq. (6.9) and relation $c \in \mathbb{N}$ imply that $e = 0$, $c = a = 1$, and $d = b = 1$. Thus $m_{2,3}^y = 3$ by Lemma 4.8, which is a contradiction to $\rho_2\rho_3\rho_2(y) = y \neq z = \rho_3\rho_2\rho_3(y)$ and (R4).

Finally, assume that $ab = 2$. If $e > 0$, then $a = 2$, $b = 1$, $c = 1$, $d = 1$, and $e = 1$ by Eq. (6.9). Then $m_{2,3}^y = 3$ by Lemma 4.8, but $\rho_2\rho_3\rho_2(y) \neq \rho_3\rho_2\rho_3(y)$, which contradicts (R4). Thus $e = 0$, and hence $c = a$, $d = b$, and $f = 0$ by Eq. (6.9) and (M2). Since $cd = 2$, we get $g = d$ by Prop. 5.1, and hence $C^x = C^y = C^z$ are Cartan matrices of type B_3 or C_3 .

Case 2: Assume that $\rho_3(x) = y$, $\rho_3(y) = x$, and $\rho_3(z) = z$. In other words, the object change diagram of $\mathcal{R}$ is

$$(6.10) \quad \begin{array}{ccccc} x & & y & & z \\ & \xrightarrow{1,3} & & \xrightarrow{2} & \\ \bullet & & \bullet & & \bullet \end{array}.$$

We apply Prop. 2.14 to $\text{Hom}(\mathcal{R})$. Let $X_x = 1_x$, $X_y = \sigma_1^x \in \text{Hom}(x, y)$, and $X_z = \sigma_2^y \sigma_1^x \in \text{Hom}(x, z)$. Then Thm. 2.6 and Prop. 2.14 imply that

$$(6.11) \quad \begin{aligned} \text{Hom}(x) \simeq \langle s_2, t, u_1, u_3 \rangle / (s_2^2, t^{m_{1,3}^x}, u_1^2, u_3^2, (u_1 s_2)^{m_{1,2}^x/3}, \\ (s_2 t^{-1} u_3 t)^{m_{2,3}^x/3}, (u_1 u_3)^{m_{1,3}^x}), \end{aligned}$$

where the inverse of the isomorphism is given by $s_2 \mapsto \sigma_2^x$, $t \mapsto \sigma_1^y \sigma_3^x$, $u_i \mapsto \sigma_1^y \sigma_2^z \sigma_i^z \sigma_2^y \sigma_1^x$ for $i \in \{1, 3\}$.

If both $m_{1,2}^x$ and $m_{2,3}^x$ are even, then

$$\text{Hom}(x)/(s_2) \simeq \langle t, u_1, u_3 \rangle / (t^{m_{1,3}^x}, u_1^2, u_3^2, (u_1 u_3)^{m_{1,3}^x}).$$

Since $m_{1,3}^x \geq 2$, the group $\text{Hom}(x)/(s_2)$ is not finite, which is a contradiction to the finiteness of $\mathcal{R}$. By symmetry and by Thm. 6.1 we may assume that $m_{1,2}^x = 3$. Then Eq. (6.11) tells that

$$(6.12) \quad \text{Hom}(x) \simeq \langle s_2, u_3, t \rangle / \langle s_2^2, u_3^2, t^{m_{1,3}^x}, (s_2 t^{-1} u_3 t)^{m_{2,3}^x/3}, (s_2 u_3)^{m_{1,3}^z} \rangle.$$

We apply Prop. 4.12 to $\text{Hom}(x)$. Since $m_{1,3}^x \geq 2$ and $m_{1,3}^z \geq 2$, we conclude that $\text{Hom}(x)$ is finite if and only if $(m_{1,3}^x, m_{2,3}^x) = (2, 3)$ or $(m_{1,3}^x, m_{2,3}^x, m_{1,3}^z) = (3, 3, 2)$. Hence $m_{2,3}^x = 3$. Using relation $m_{1,2}^x = 3$, Lemma 4.8, (C2), and Thm. 6.1, we obtain that

$$C^x = \begin{pmatrix} 2 & -1 & -a \\ -1 & 2 & -1 \\ -b & -1 & 2 \end{pmatrix}, C^y = \begin{pmatrix} 2 & -1 & -a \\ -1 & 2 & -1 \\ -b & -1 & 2 \end{pmatrix}, C^z = \begin{pmatrix} 2 & -1 & -c \\ -1 & 2 & -1 \\ -d & -1 & 2 \end{pmatrix},$$

where $a, b, c, d \in \mathbb{N}_0$. If $m_{1,3}^x = 3$ and $m_{1,3}^z = 2$ then $c = d = 0$ and $a = b = 1$ by Lemma 4.8. Then $a + 1 \neq c + 1$, which is a contradiction to Lemma 4.9. If $m_{1,3}^x = 2$ then $a = b = 0$, and hence $c = d = 0$ by Lemma 4.9 and (M2). Then $C^x = C^y = C^z$ are Cartan matrices of type A_3 . This proves the theorem in Case 2.

Case 3: Assume that $\rho_3(x) = z$, $\rho_3(y) = y$, and $\rho_3(z) = x$. For all $a \in A$ and $i, j \in \{1, 2, 3\}$ with $i \neq j$ the relations $(\rho_i \rho_j)^3(a) = a$ and $\rho_i \rho_j(a) \neq a$ hold, and hence $3 | m_{i,j}^a$ by (R4). In particular, $c_{ij}^a < 0$ for all $a \in A$ and $i, j \in \{1, 2, 3\}$ with $i \neq j$.

Calculate $\sigma := \sigma_2^x \sigma_3^z \sigma_2^y \sigma_1^x \in \text{Hom}(x)$. Since $\mathcal{R}$ is finite, σ has finite order, and hence $\sigma = \text{id}$ or $\text{tr } \sigma < 3$. Direct calculation shows that $\sigma(\alpha_1) \in -\alpha_1 + \mathbb{Z}\alpha_2 + \mathbb{Z}\alpha_3$ and

$$\begin{aligned} \text{tr } \sigma &= c_{12}^x c_{21}^x - c_{12}^x c_{23}^x c_{31}^z + (c_{12}^x c_{21}^y - 1)(c_{23}^x c_{32}^z - 1) \\ &\quad + c_{13}^x c_{31}^z - c_{13}^x c_{21}^y c_{32}^z + c_{23}^y c_{32}^z - 2. \end{aligned}$$

By the above conclusion one obtains that $\text{tr } \sigma \geq 3$ and $\sigma \neq \text{id}$, which is a contradiction. Thus there are no finite root systems of type $\mathcal{C}$ in this case. This finishes the proof of the theorem. $\square$

For the classification of root systems of type $\mathcal{C}$ of rank higher than 3 the following proposition will be useful.

Proposition 6.4. *Let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a connected Cartan scheme with $I = \{1, 2, 3, 4\}$ and three objects x, y, z , and let $\mathcal{R}$ be a finite root system of type $\mathcal{C}$. Assume that $\rho_1(x) = y$, $\rho_2(y) = z$, and $c_{12}^a = c_{21}^a = -1$ for all $a \in \{x, y, z\}$. Then $\mathcal{R}$ is standard.*

Proof. Step 1. Both $\mathcal{R}|_{\{1,2,3\}}$ and $\mathcal{R}|_{\{1,2,4\}}$ are standard.

Since $\mathcal{R}|_{\{1,2\}}$ is connected and irreducible, either $\mathcal{R}|_{\{1,2,3\}} = \mathcal{R}|_{\{1,2\}} \oplus \mathcal{R}|_{\{3\}}$, or $\mathcal{R}|_{\{1,2,3\}}$ is connected and irreducible. Then $\mathcal{R}|_{\{1,2,3\}}$ is standard — in the first case, since $\mathcal{R}|_{\{1,2\}}$ is standard, and in the second case by Thm. 6.3. Similarly, $\mathcal{R}|_{\{1,2,4\}}$ is standard.

Step 2. If $\rho_3(x) \neq x$ then $\mathcal{R}$ is standard.

By Thm. 6.1, $\mathcal{R}|_{\{2,3\}}$ is connected and irreducible. Hence $\mathcal{R}|_{\{1,2,3\}}$ is connected and irreducible, and Thm. 6.3 implies that $\rho_3(x) = y$ and $c_{23}^a = c_{32}^a = -1$ for all $a \in \{x, y, z\}$. Thus Step 1 gives that $\mathcal{R}|_{\{1,2,3\}}$, $\mathcal{R}|_{\{1,2,4\}}$, and $\mathcal{R}|_{\{3,2,4\}}$ are standard, hence $\mathcal{R}$ is standard, see Rem. 4.2.

Step 3. $\mathcal{R}$ is standard.

If $\rho_3 \neq \text{id}$ or $\rho_4 \neq \text{id}$, then $\mathcal{R}$ is standard by Step 2. Assume now that $\rho_3(a) = \rho_4(a) = a$ for all $a \in A$. If $\mathcal{R} = \mathcal{R}|_{\{1,2\}} \oplus \mathcal{R}|_{\{3,4\}}$, then $c_{ij}^y = 0$ for all $i \in \{1, 2\}$, $j \in \{3, 4\}$. Thus $c_{jl}^x = c_{jl}^y = c_{jl}^z$ for all $j \in \{3, 4\}$ and $l \in I$ by Lemma 4.5. Since $\mathcal{R}|_{\{1,2,3\}}$ and $\mathcal{R}|_{\{1,2,4\}}$ are standard by Step 1, it follows that $c_{jl}^x = c_{jl}^y = c_{jl}^z$ for all $j \in \{1, 2\}$ and $l \in I$. Hence $\mathcal{R}$ is standard.

Assume now that $\mathcal{R}|_{\{1,2,3\}}$ is irreducible. By Thm. 6.3 we may assume that $c_{13}^a = c_{31}^a = 0$, $c_{23}^a c_{32}^a = 2$, and $m_{2,3}^a = 4$ for all $a \in \{x, y, z\}$. Then Thm. 2.6 and Prop. 2.14 give that

$$(6.13) \quad \begin{aligned} \text{Hom}(x) \simeq \langle s_2, s_3, s_4, t_3, t_4, u_1, u_3, u_4 \rangle / (s_2^2, s_3^2, s_4^2, t_3^2, t_4^2, u_1^2, \\ u_3^2, u_4^2, u_1 s_2, s_3 t_3, (s_4 t_4)^{m_{1,4}^x/2}, (s_2 s_3)^4, \\ (s_2 s_4)^{m_{2,4}^x}, (s_3 s_4)^{m_{3,4}^x}, (t_3 u_3)^2, (t_4 u_4)^{m_{2,4}^y/2}, \\ (t_3 t_4)^{m_{3,4}^y}, (u_1 u_3)^2, (u_1 u_4)^{m_{1,4}^z}, (u_3 u_4)^{m_{3,4}^z}), \end{aligned}$$

where the inverse of the isomorphism is given by the map $s_i \mapsto \sigma_i^x$ for all $i \in \{2, 3, 4\}$, $t_i \mapsto \sigma_1^y \sigma_i^y \sigma_1^x$ for all $i \in \{3, 4\}$, and $u_i \mapsto \sigma_1^y \sigma_2^z \sigma_i^z \sigma_2^y \sigma_1^x$ for all $i \in \{1, 3, 4\}$. Therefore

$$(6.14) \quad \begin{aligned} \text{Hom}(x) \simeq \langle s_2, s_3, s_4, t_4, u_3, u_4 \rangle / (s_2^2, s_3^2, s_4^2, t_4^2, u_3^2, u_4^2, \\ (s_4 t_4)^{m_{1,4}^x/2}, (s_2 s_3)^4, (s_2 s_4)^{m_{2,4}^x}, (s_3 s_4)^{m_{3,4}^x}, (s_3 u_3)^2, \\ (t_4 u_4)^{m_{2,4}^y/2}, (s_3 t_4)^{m_{3,4}^y}, (s_2 u_3)^2, (s_2 u_4)^{m_{1,4}^z}, (u_3 u_4)^{m_{3,4}^z}). \end{aligned}$$

If $m_{1,4}^x > 2$ and $m_{2,4}^y > 2$, then $\text{Hom}(x)$ is a Coxeter group without Coxeter relation between s_2 and t_4 , which is a contradiction to the finiteness of $\mathcal{R}$.

Assume first that $m_{1,4}^x = 2$. Since $m_{1,3}^x = 2$ as well, we obtain that $c_{13}^x = c_{14}^x = 0$, and hence $m_{3,4}^x = m_{3,4}^y$. Then

$$(6.15) \quad \begin{aligned} \text{Hom}(x) \simeq & \langle s_2, s_3, s_4, u_3, u_4 \rangle / (s_2^2, s_3^2, s_4^2, u_3^2, u_4^2, (s_2 s_3)^4, \\ & (s_2 s_4)^{m_{2,4}^x}, (s_3 s_4)^{m_{3,4}^x}, (s_3 u_3)^2, (s_4 u_4)^{m_{2,4}^y/2}, \\ & (s_2 u_3)^2, (s_2 u_4)^{m_{1,4}^z}, (u_3 u_4)^{m_{3,4}^z}). \end{aligned}$$

If $m_{2,4}^y > 2$, then there is no Coxeter relation between s_4 and u_3 , which is a contradiction, and hence $m_{2,4}^y = 2$.

Similarly, if $m_{1,4}^x > 2$ and $m_{2,4}^y = 2$, then there is no Coxeter relation between s_4 and u_3 .

We are left with the case $m_{1,4}^x = m_{2,4}^y = 2$. Then relations $c_{14}^x = c_{42}^y = 0$ and Lemma 4.5 imply that $c_{42}^x = c_{42}^y = 0$, and hence $m_{2,4}^x = 2$. Similarly, $c_{24}^y = 0$ implies that $c_{41}^z = c_{41}^y = c_{41}^x = 0$, hence $m_{1,4}^z = 2$. Therefore

$$(6.16) \quad \begin{aligned} \text{Hom}(x) \simeq & \langle s_2, s_3, s_4, u_3 \rangle / (s_2^2, s_3^2, s_4^2, u_3^2, (s_2 s_3)^4, (s_2 s_4)^2, \\ & (s_3 s_4)^{m_{3,4}^x}, (s_3 u_3)^2, (s_2 u_3)^2, (u_3 s_4)^{m_{3,4}^z}). \end{aligned}$$

The finiteness of $\text{Hom}(x)$ implies that $m_{3,4}^x, m_{3,4}^z \in \{2, 3\}$. If $m_{3,4}^x = 2$, then $m_{1,4}^x = m_{2,4}^x = 0$ implies that $\mathcal{R} = \mathcal{R}|_{\{1,2,3\}} \oplus \mathcal{R}|_{\{4\}}$, and hence $\mathcal{R}$ is standard, since $\mathcal{R}|_{\{1,2,3\}}$ is standard. Further, $m_{3,4}^z = 2$. Similarly, the assumption $m_{3,4}^z = 2$ leads to the same result.

Suppose now that $m_{3,4}^x = m_{3,4}^z = 3$, that is, $c_{34}^x = c_{43}^x = c_{34}^z = c_{43}^z = -1$ by Lemma 4.8. Since $c_{13}^x = c_{14}^x = 0$, Lemma 4.5 implies that $c_{34}^y = c_{43}^y = -1$, that is, $C^x = C^y = C^z$ is a Cartan matrix of type F_4 . This proves the proposition. $\square$

Theorem 6.5. *Let $\theta \in \mathbb{N}$ with $\theta \geq 4$, let $\mathcal{C} = \mathcal{C}(I, A, (\rho_i)_{i \in I}, (C^a)_{a \in A})$ be a connected Cartan scheme with $I = \{1, 2, \dots, \theta\}$ and with 3 objects, and let $\mathcal{R}$ be a finite irreducible root system of type $\mathcal{C}$. Then $\mathcal{R}$ is standard, and the corresponding Cartan matrix C is indecomposable and of type B_4 , C_4 , D_4 , or F_4 . The object change diagram of $\mathcal{R}$ can be chosen to be*

$$\begin{array}{ll} \bullet \xrightarrow{1,3} \bullet \xrightarrow{2} \bullet & \text{if } C \text{ is of type } B_4 \text{ or } C_4, \\ \bullet \xrightarrow{1,3,4} \bullet \xrightarrow{2} \bullet & \text{if } C \text{ is of type } D_4, \text{ and} \\ \bullet \xrightarrow{1} \bullet \xrightarrow{2} \bullet & \text{if } C \text{ is of type } F_4. \end{array}$$

Proof. As explained at the beginning of this section, we can choose $x, y, z \in A$ and use a permutation of I such that the object change diagram of $\mathcal{R}|_{\{1,2\}}$ is the one in Eq. (6.1). Since $\mathcal{R}$ is irreducible, we

can assume that $\mathcal{R}|_{\{1,2,3\}}$ is irreducible. Then Thm. 6.3 gives that $\mathcal{R}|_{\{1,2,3\}}$ is standard and $m_{1,2}^x \in \{2, 3, 4\}$. Further, since $\rho_2\rho_1(x) = z \neq y = \rho_1\rho_2(x)$ and $\rho_1\rho_2\rho_1(x) = z = \rho_2\rho_1\rho_2(x)$, (R4) yields that $m_{1,2}^x = 3$. Hence $c_{12}^a = c_{21}^a = -1$ for all $a \in A$ by Lemma 4.8. By Prop. 6.4 we obtain that $\mathcal{R}|_{\{1,2,i,j\}}$ is standard for all $i, j \in I \setminus \{1, 2\}$ with $i \neq j$. Hence $\mathcal{R}$ is standard by Rem. 4.2.

Finally, Thm. 6.3 and elementary argumentations with Dynkin diagrams imply that there is no connected irreducible standard root system, where the Cartan matrices are of type A_4 or of rank bigger than 4, and that connected irreducible standard root systems of rank 4 have the given object change diagrams. The rigorous proof is left to the reader. $\square$

7. APPENDIX

In the following table we list all non-standard connected irreducible finite Weyl groupoids with at most three objects. The last column contains the stabilizer group of an object. It is a Coxeter group of the indicated type.

$\mathcal{W}$	$ A $	$ I $	$ \mathcal{W} $	$ R_+^a $	$\text{Hom}(a)$
Thm. 5.4(2)	2	2	32	8	B_2
Thm. 5.4(2)	2	2	48	12	G_2
Thm. 5.4(3)	2	3	192	13	B_3
Thm. 5.4(4)	2	3	192	13	B_3
Thm. 6.1(3)	3	2	36	6	$A_1 \times A_1$
Thm. 6.1(4)	3	2	72	12	B_2
Thm. 6.1(5)	3	2	72	12	B_2
Thm. 6.1(6)	3	2	108	18	G_2
Thm. 6.1(7)	3	2	108	18	G_2

REFERENCES

- [AHS08] N. Andruskiewitsch, I. Heckenberger, and H.-J. Schneider, *The Nichols algebra of a semisimple Yetter-Drinfeld module*, Preprint arXiv:0803.2430 (2008), 52 pages.
- [AS98] N. Andruskiewitsch and H.-J. Schneider, *Lifting of quantum linear spaces and pointed Hopf algebras of order p^3* , J. Algebra **209** (1998), 658–691.
- [AS00] ———, *Finite quantum groups and Cartan matrices*, Adv. Math. **154** (2000), 1–45.
- [BMM99] M. Broué, G. Malle, and J. Michel, *Towards spetses. I*, Transform. Groups **4** (1999), no. 2-3, 157–218, Dedicated to the memory of Claude Chevalley.
- [Bou68] N. Bourbaki, *Groupes et algèbres de Lie, ch. 4, 5 et 6*, Éléments de mathématique, Hermann, Paris, 1968.

- [Hec05] I. Heckenberger, *Classification of arithmetic root systems of rank 3*, Actas del “XVI Coloquio Latinoamericano de Álgebra” (Colonia, Uruguay, 2005), 2005, pp. 227–252.
- [Hec06a] ———, *Classification of arithmetic root systems*, Preprint math.QA/0605795 (2006), 67 pages.
- [Hec06b] ———, *The Weyl groupoid of a Nichols algebra of diagonal type*, Invent. Math. **164** (2006), 175–188.
- [Hec07] ———, *Examples of finite-dimensional rank 2 Nichols algebras of diagonal type*, Compositio Math. **143** (2007), 165–190.
- [HY08] I. Heckenberger and H. Yamane, *A generalization of Coxeter groups, root systems, and Matsumoto’s theorem*, Math. Z. **259** (2008), 255–276.
- [Kac90] V.G. Kac, *Infinite dimensional Lie algebras*, Cambridge Univ. Press, 1990.

MICHAEL CUNTZ, FACHBEREICH MATHEMATIK, UNIVERSITÄT KAISERSLAUTERN, POSTFACH 3049, D-67653 KAISERSLAUTERN, GERMANY

E-mail address: cuntz@mathematik.uni-kl.de

ISTVÁN HECKENBERGER, MATHEMATISCHES INSTITUT, LUDWIG-MAXIMILIANS-UNIVERSITÄT MÜNCHEN, THERESIENSTR. 39, D-80333 MÜNCHEN, GERMANY

E-mail address: i.heckenberger@gmail.com